

SITE SYNOPSIS

SITE NAME: SLANEY RIVER VALLEY

SITE CODE: 000781

This site comprises the freshwater stretches of the Slaney as far as the Wicklow Mountains; a number of tributaries the larger of which include the Bann, Boro, Glasha, Clody, Derry, Derreen, Douglas and Carrigower Rivers; the estuary at Ferrycarrig and Wexford Harbour. The site flows through the counties of Wicklow, Wexford and Carlow. Towns along the site but not in it are Baltinglass, Hacketstown, Tinahely, Tullow, Bunclody, Camolin, Enniscorthy and Wexford. The river is up to 100 m wide in places and is tidal at the southern end from Edermine Bridge below Enniscorthy. In the upper and central regions almost as far as the confluence with the Derry River the geology consists of granite. Above Kilcarr Bridge, the Slaney has cut a gorge into the granite plain. The Derry and Bann Rivers are bounded by a narrow line of uplands which corresponds to schist outcrops. Where these tributaries cut through this belt of hard rocks they have carved deep gorges, more than two miles long at Tinahely and Shillelagh. South of Kildavin the Slaney flows through an area of Ordovician slates and grits.

The site is a candidate SAC selected for alluvial wet woodlands, a priority habitat on Annex I of the E.U. Habitats Directive. The site is also selected as a candidate SAC for floating river vegetation, estuaries, tidal mudflats and old oak woodlands, all habitats listed on Annex I of the E.U. Habitats Directive. The site is further selected for the following species listed on Annex II of the same directive - Sea Lamprey, River Lamprey, Brook Lamprey, Freshwater Pearl Mussel, Twaite Shad, Atlantic Salmon and Otter.

Floating river vegetation is found along much of the freshwater stretches within the site. Species present here include Pond Water-crowfoot (*Ranunculus peltatus*), Water-crowfoot (*Ranunculus* spp.), Canadian Pondweed (*Elodea canadensis*), Broad-leaved Pondweed (*Potamogeton natans*), Water Milfoil (*Myriophyllum* spp.), Common Club-rush (*Scirpus lacustris*), Water-starwort (*Callitriche* spp.), Hemlock Water-dropwort, Fine-leaved Water-dropwort (*Oenanthe aquatica*), Common Duckweed (*Lemna minor*), Yellow Water-lily (*Nuphar lutea*), Unbranched Bur-reed (*Sparganium emersum*) and the moss *Fontinalis antipyretica*. Two rare aquatic plant species have been recorded in this site: Short-leaved Water-starwort (*Callitriche truncata*), a very rare, small aquatic herb found nowhere else in Ireland; and Opposite-leaved Pondweed (*Groenlandia densa*), a species that is legally protected under the Flora Protection Order, 1999.

Good examples of wet woodland are found associated with Macmine marshes, along banks of the Slaney and its tributaries and within reed swamps. Grey Willow (*Salix cinerea*) scrub and pockets of wet woodland dominated by Alder (*Alnus glutinosa*) have become established in places. Ash (*Fraxinus excelsior*) and Birch (*Betula pubescens*) are common in the latter and the ground flora is typical of wet woodland with Meadowsweet (*Filipendula ulmaria*), Angelica (*Angelica sylvestris*), Yellow Iris, Horsetail (*Equisetum* spp.) and occasional tussocks of Greater Tussock-sedge (*Carex paniculata*). These woodlands have been described as two types: one is quite eutrophic, is dominated by Willow and is subject to a tidal influence. The

other is flushed or spring-fed subject to waterlogging but not to flooding and is dominated by Alder and Ash.

Old oak woodlands are best represented at Tomnafinnoge though patches are present throughout the site. At Tomnafinnoge the wood is dominated by mature, widely spaced Sessile Oak (*Quercus petraea*), which were planted around 1700, with some further planting in 1810. There is now a varied age structure with overmature, mature and young trees; the open canopy permits light to reach the forest floor and encourages natural regeneration of Oak. As well as Oak, the wood includes the occasional Beech (*Fagus sylvatica*), Birch (*Betula* sp.), Rowan (*Sorbus aucuparia*) and Scots Pine (*Pinus sylvestris*).

The shrub layer is well-developed with Hazel (*Corylus avellana*) and Holly (*Ilex aquifolium*) occurring. The ground layer consists of Great Wood-rush (*Luzula sylvatica*) and Bilberry (*Vaccinium myrtillus*), with some Bracken (*Pteridium aquilinum*) and Brambles (*Rubus fruticosus* agg.). Herbaceous species in the ground layer include Primrose (*Primula vulgaris*), Wood-sorrel (*Oxalis acetosella*), Common Cow-wheat (*Melampyrum pratense*) and Bluebell (*Hyacinthoides non-scripta*). Many of the trees carry an epiphytic flora of mosses, Polypody Fern (*Polypodium vulgare*), and lichens such as *Usnea comosa*, *Evernia prunastri*, *Ramalina* spp. and *Parmelia* spp.

Tomnafinnoge Wood is a remnant of the ancient Shillelagh Oak woods, and it appears that woodland has always been present on the site. In the past, the wood was managed as a Hazel coppice with Oak standards a common form of woodland management in England but not widely practised in Ireland. The importance of the woodland lies in the size of the trees, their capacity to regenerate, their genetic continuity with ancient woodland and their historic interest. The nearest comparable stands are at Abbeyleix, Co. Laois and Portlaw, Co. Waterford.

Below Enniscorthy there are several areas of woodland with a mixed canopy of Oak, Beech, Sycamore (*Acer pseudoplatanus*), Ash and generally a good diverse ground flora. Near the mouth of the river at Ferrycarrig is a steep south facing slope covered with Oak woodland. Holly and Hazel are the main species in the shrub layer and a species-rich ground flora typical of this type of Oak woodland has abundant ferns - *Dryopteris filix-mas*, *Polystichum setiferum*, *Phyllitis scolopendrium* - and mosses - *Thuidium tamariscinum*, *Mnium hornum*, *Eurynchium praelongum*.

North of Bunclody, the river valley still has a number of dry woodlands though these have mostly been managed by the estates with the introduction of Beech and occasional conifers. The steeper sides are covered in a thick scrub from which taller trees protrude. At the southern end of the site, the Red Data Book species Yellow Archangel (*Lamiastrum galeobdolon*) occurs. Three more Red Data Book species have also been recorded from the site: Basil Thyme (*Acinos arvensis*), Blue Fleabane (*Erigeron acer*) and Small Cudweed (*Filago minima*). A nationally rare species Summer Snowflake (*Leucojum aestivum*) is also found within the site.

Mixed woodlands occur at Carrickduff and Coolaphuca in Bunclody. Oak trees, which make up the greater part of the canopy, were originally planted and at the present time are not

regenerating actively. In time, if permitted, the woodland will probably go to Beech. A fair number of Yew (*Taxus baccata*) trees have also reached a large size and these, together with Holly give to the site the aspect of a south-western Oak wood.

The site is considered to contain a very good example of the extreme upper reaches of an estuary. Tidal reedbeds with wet woodland are present in places. The fringing reed communities support Sea Club-rush (*Scirpus maritimus*), Grey Club-rush (*S. tabernaemontani*) and abundant Common Reed (*Phragmites australis*). Other species occurring are Bulrush (*Typha latifolia*), Reed Canary-grass (*Phalaris arundinacea*) and Branched Bur-reed (*Sparganium erectum*). The reed-swamp is extensive around Macmine, where the river widens and there are islands with swamp and marsh vegetation.

Further south of Macmine are expanses of intertidal mudflats and sandflats and shingly shore often fringed with a narrow band of salt marsh and brackish vegetation. Narrow shingle beaches up to 10 m wide occur in places along the river banks and are exposed at low tide. Upslope the shingle is sometimes colonised by Saltmarsh Rush (*Juncus gerardi*), Townsend's Cord-grass (*Spartina townsendii*), Common Saltmarsh-grass (*Puccinellia maritima*), Sea Aster (*Aster tripolium*), Hemlock Water-dropwort (*Oenanthe crocata*) and Himalayan Balsam (*Impatiens glandulifera*).

Wexford Harbour is an extensive, shallow estuary which dries out considerably at low tide exposing large expanses of mudflats and sandflats. The harbour is largely sheltered by the Raven Point to the north and Rosslare Point in the south.

Other habitats present within the site include species-rich marsh in which sedges such as *Carex disticha*, *Carex riparia* and *Carex vesicaria* are common. Among the other species found in this habitat are Yellow Iris (*Iris pseudacorus*), Water Mint (*Mentha aquatica*), Purple Loosestrife (*Lythrum salicaria*) and Soft Rush (*Juncus effusus*). Extensive marshes occur to the west of Castlebridge associated with the tidal areas of the River Sow.

The site supports populations of several species listed on Annex II of the EU Habitats Directive including the three Lampreys - Sea Lamprey (*Petromyzon marinus*), River Lamprey (*Lampetra fluviatilis*) and Brook Lamprey (*Lampetra planeri*), Otter (*Lutra lutra*), Salmon (*Salmo salar*), small numbers of Freshwater Pearl Mussel (*Margaritifera margaritifera*) and in the tidal stretches, Twait Shad (*Alosa fallax fallax*). A survey of the Derreen River in 1995 estimated the population of Freshwater Pearl Mussel at about 3,000 individuals. This is a significant population, especially in the context of eastern Ireland. The Slaney is primarily a spring salmon fishery and is regarded as one of the top rivers in Ireland for early spring fishing. The upper Slaney and tributary headwaters are very important for spawning.

The site supports important numbers of birds in winter. Little Egret are found annually along the river. This bird is only now beginning to gain a foothold in Ireland and the south-east appears to be its stronghold. Nationally important numbers of Black-tailed Godwit, Teal, Tufted Duck, Mute Swan, Little Grebe and Black-headed Gull are found along the estuarine stretch of the river. The mean of the maximum counts over four winters (1994/98) along the stretch between Enniscorthy and Ferrycarrig is: Little Egret (6), Golden Plover (6), Wigeon (139), Teal (429), Mallard (265), Tufted Duck (171), Lapwing (603), Shelduck (16), Black-tailed Godwit (93), Curlew (81), Red-breasted Merganser (11), Black-headed Gull (3030), Goldeneye (45), Oystercatcher (19), Redshank (65), Lesser Black-backed Gull (727), Herring

Gull (179), Common Gull (67), Grey Heron (39), Mute Swan (259) and Little Grebe (17). Wexford Harbour provides extensive feeding grounds for wading birds and Little Terns, which are listed on Annex I of the E.U. Birds Directive have bred here in the past.

The Reed Warbler, which is a scarce breeding species in Ireland, is regularly found in Macmine Marshes but it is not known whether or not it breeds in the site. The Dipper also occurs on the river. This is a declining species nationally.

The site supports many of the mammal species occurring in Ireland. Those which are listed in the Irish Red Data Book include Pine Marten, Badger, Irish Hare and Daubenton's Bat. Common Frog (*Rana temporaria*), another Red Data Book species, also occurs within the site.

Agriculture is the main landuse. Arable crops are important. Improved grassland and silage account for much of the remainder. The spreading of slurry and fertiliser poses a threat to the water quality of this salmonid river and to the populations of Annex II animal species within it. Run-off is undoubtedly occurring, as some of the fields slope steeply directly to the river bank. In addition, cattle have access to the site in places. Fishing is a main tourist attraction along stretches of the Slaney and its tributaries and there are a number of Angler Associations, some with a number of beats. Fishing stands and styles have been erected in places. Both commercial and leisure fishing takes place. There are some gravel pits along the river below Bunclody and many of these are active. There is a large landfill site adjacent to the river close to Hacketstown and at Killurin. Boating, bait-digging and fishing occur in parts of Wexford Harbour.

Waste water outflows, runoff from intensive agricultural enterprises, a meat factory at Clohamon and a landfill site adjacent to the river and further industrial development upstream in Enniscorthy and in other towns could all have potential adverse impacts on the water quality unless they are carefully managed. The spread of exotic species is reducing the quality of the woodlands.

The site supports populations of several species listed on Annex II of the EU Habitats Directive, and habitats listed on Annex I of this directive, as well as important numbers of wintering wildfowl including some species listed on Annex I of the EU Birds Directive. The presence of wet and broad-leaved woodlands increases the overall habitat diversity and the occurrence of a number of Red Data Book plant and animal species adds further importance to the Slaney River site.

07.12.2005