

Number 527552

# Certificate of Incorporation

I hereby certify that

**STARRUS ECO HOLDINGS LIMITED**

is this day incorporated under  
the Companies Acts 1963 to 2012,  
and that the company is limited.

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

Given under my hand at Dublin, this  
**Wednesday, the 15th day of May, 2013**



for Registrar of Companies



# CLLIB HOLDINGS.


## DOCK ROAD LIMERICK.

Mr Malcolm Dowling  
Group Environmental Compliance Manager  
Greenstar  
Unit 6 Ballyogan Business Park  
Ballyogan Road,  
Dublin 18

Dear Mr Dowling,


Starrus Eco Holdings Ltd T/A Greenstar who is the leaseholder of the lands at Ballykeeffe, Dock Road, County Limerick, intends to apply for a Licence to increase the amount of waste accepted at the facility to 130,000 tonnes annually. We understand that the proposed increase in the amount of wastes accepted does not require the construction of any new buildings or the alteration to the existing infrastructure. We give our consent to Greenstar to make the applications.

Signed:




Date: 20 03 2014

YOURS SINCERELY,



FOR AND ON BEHALF OF

CLLIB HOLDINGS.



- SITE BOUNDARY (RED)
- SURFACEWATER DRAINAGE
- FOULWATER DRAINAGE
- NOISE MONITORING LOCATION
- ⊕ DUST MONITORING LOCATION
- ⊠ GROUNDWATER MONITORING LOCATION
- + FOULWATER MONITORING LOCATION

NOTES

Monitoring locations  
Grid references  
Eastings Northings

DM-1	- 154913, 155617
DM-2	- 154842, 155714
DM-3	- 155003, 155750
Ni-1	- 154840, 155712
Ni-2	- 154920, 155745
Ni-3	- 154910, 155618
FE-1A	- 154974, 155769
WS-9	- 155003, 155750
WS-10	- 154922, 155776
FE-2A	- 154858, 155720
GW-M1	- 154923, 155667
GW-M2	- 154920, 155770
GW-M3	- 155049, 155684

For information purposes only.  
Consent of copyright owner required for any other use.

A	27/01/13	PRELIMINARY ISSUE	MW	SM	**
REV	DATE	DESCRIPTION	DRN	CHKD	APP

**O'Callaghan Moran & Associates**  
Granary House, Rutland Street,  
Cork, Ireland.  
Tel: (021) 4321521 Fax: (021) 4321522  
email: ocm@mdigo.ie


This drawing is the property of O'Callaghan Moran & Associates and shall not be used, reproduced or disclosed to anyone without the prior written permission of O'Callaghan Moran & Associates and shall be returned upon request.

CLIENT  
**Greenstar**

TITLE  
**Emission & Monitoring  
LIMERICK**


SCALE	DRAWING No.	REV.
1:1000 A3	01	A

DOS Filename : \*\*\*\*\*



NOTES

REV	DATE	DESCRIPTION	DRN	CHKD	APP
A	27/01/13	PRELIMINARY ISSUE	MW	SM	**


**O'Callaghan Moran & Associates.**  
 Granary House, Rutland Street,  
 Cork, Ireland.  
 Tel: (021) 4321521 Fax: (021) 4321522  
 email: ocm@indigo.ie

This drawing is the property of O'Callaghan Moran & Associates and shall not be used, reproduced or disclosed to anyone without the prior written permission of O'Callaghan Moran & Associates and shall be returned upon request.

CLIENT  
**Greenstar E.S Ltd**

TITLE  
**NOTICE LOCATIONS**

SCALE 1:1000 A3	DRAWING No. B2	REV. A
-----------------------	-------------------	-----------

DOS Filename : \*\*\*\*\*

## **B6 Planning History**

In 1994, the landowner Cussen & Co Crane Hire Limited, now CLIIB obtained planning permission (876/94) for the retention of a workshop extension, vehicle wash and compound and the erection of 2 No. 5,000 gallon fuel tanks with pumps and security fencing.

In 1995, Cussen obtained planning permission (162/95) for the erection of 6 No. industrial units and the provision of a wastewater treatment plant. Cussen also obtained planning permission (968/95) for the retention of raised lands.

In 2000 Cussen was granted planning permission (PL 13.110811) for upgrade works, which included the construction of Building 1 and ancillary works.

IPODEC Ireland Ltd. which was renamed Onyx and subsequently Veolia Environmental Services Ireland Ltd (VESI), acquired the Cussen waste business in 2001. In October 2002, VESI was granted planning permission (02/984) for the construction of Building 2 and ancillary works and these were completed in 2003.

On 13<sup>th</sup> March 2014 Limerick County Council granted permission to increase the annual tonnage to 130,000 tonnes (13/300).

For inspection purposes only.  
Consent of copyright owner required for any other use.

**LIMERICK COUNTY COUNCIL**

**PLANNING AND DEVELOPMENT ACTS 2000-2013**

**NOTIFICATION OF DECISION TO GRANT**

Greenstar Environmental Services Ltd (In Receivership)  
c/o O'Callaghan Moran & Associates  
Granary House  
Rutland Street  
Cork

**Planning Register Number:** 13/300  
**Valid Application Received:** 22/05/2013  
**Further Information Received Date:** 17/01/2014

In pursuance of the powers conferred upon them by the above-mentioned Act, Limerick County Council has by Order dated **13<sup>th</sup> March, 2014** decided for the reason set out in the First Schedule hereto, to GRANT **PERMISSION** for development of land in accordance with the documents submitted namely:-**an increase in the amount of waste accepted annually to 130,000 tonnes. The proposed increase does not require the construction/provision of any new buildings or structures (The development will require a revision of the Waste Licence granted by the Environmental Protected Agency, also, this application is accompanied by an Environmental Impact Statement (EIS)) at existing Materials Recovery Facility Ballykeeffe Dock Road** subject to the 5 condition(s) and the reasons for the imposition of the said condition(s) as set out in the Second Schedule.

Signed on behalf of said Council \_\_\_\_\_  
for **DIRECTOR OF SERVICES**  
**ECONOMIC DEVELOPMENT & PLANNING**

Date: **13<sup>th</sup> March, 2014**

Under Article 20 of the Planning & Development Regulations 2001 - 2013 the applicant shall remove the site notice in respect of the application following notification of the Planning Authority's decision. In deciding the planning application, the Planning Authority, in accordance with Section 34(3) of the Planning & Development Acts 2000 – 2013 has had regard to submissions/observations received (if any) in accordance with the Planning & Development Regulations 2001 – 2013.

If there is no appeal to An Bord Pleanála a grant of permission shall be issued as soon as may be but not earlier than 3 working days after the expiration of the period for making of an appeal (see footnote).

**THIS NOTICE IS NOT A GRANT OF PERMISSION AND WORK SHOULD NOT COMMENCE UNTIL PLANNING PERMISSION IS GRANTED.**

**NOTE:**

An appeal against a decision of a planning authority under the provisions of the Planning & Development Acts, 2000-2013 may be made to An Bord Pleanála at any time before the expiration of the appropriate period and on payment of the appropriate fee, by an applicant for permission or any person who made submissions or observations in writing in relation to the planning application. An appeal by a person who made submissions or observations must be accompanied by the acknowledgement of receipt of the submissions or observations from the planning authority. Any such appeal must be made in writing and received by the Board within 4 weeks beginning on the date of the making of the decision by the planning authority. The appeal must be fully complete from the start otherwise it will be invalid. It is very important to note that any appeal referrals under the 2000 to 2013 Planning & Development Acts which are not accompanied by the correct fee will be invalid.

/P.T.O.

The scale of fees payable to An Bord Pleanála in respect of appeals is set out hereunder:

Case Type	Appeal received on or after 5 <sup>th</sup> September 2011
<b>Planning Acts</b>	
a. Application for strategic infrastructure development or a request to alter the terms of such development already permitted or approved.	€100,000
b. Appeal against a decision of a planning authority on a planning application relating to commercial development, made by the person by whom the planning application was made, where the application included retention of development.	€4,500 or €9,000 if *EIS or **NIS involved
c. Appeal against a decision of a planning authority on a planning application relating to commercial development, made by the person by whom the planning application was made, other than an appeal mentioned at (b).	€1,500 or €3,000 if *EIS or **NIS involved
d. Appeal against a decision of a planning authority on a planning application made by the person by whom the planning application was made, where the application relates to retention of development, other than an appeal mentioned at (b) or (c) (non-commercial development).	€660
e. 1 <sup>st</sup> party appeal solely against contribution condition(s) – (2000 Act Section 48 or 49).	€220
f. Appeal other than an appeal mentioned at (b), (c), (d) or (h)	€220
g. Application for leave to appeal.	€110
h. Appeal following a grant of leave to appeal.	€110
i. Referral.	€220
j. Reduced fee (payable by specified bodies).	€110
k. Submissions or observations (by observer) on strategic infrastructure development applications, appeals and referrals.	€50
l. Request from a party for an oral hearing.	€50
*EIS - Environmental Impact Statement	
**NIS - Natura Impact Statement	

Submissions or observations on appeals made by third parties must be received by the Board within 4 weeks from the receipt of the appeal by the Board and the fee in this case is €50. Development consisting of the provision of two or more dwellings is classed as commercial development for the purposes of an appeal.

Should you wish to make an appeal, the following documents are available on [www.pleanala.ie](http://www.pleanala.ie)

- A Planning Appeal Form/Checklist and
- A Guide to making a Planning Appeal.

Appeals should be addressed to An Bord Pleanála, 64 Marlborough Street, Dublin 1.

**-PLANNING REGISTER REFERENCE NUMBER: 13/300**

**FIRST SCHEDULE**

Having regard to the nature of the proposed development, it is considered that subject to compliance with the conditions as set out in the Second Schedule, the proposed development would be in accordance with the proper planning and sustainable development of the area.

**SECOND SCHEDULE**

1. The development shall be carried out in accordance with the plans and particulars lodged with the application and the E.I.S. on the 22<sup>nd</sup> May 2013 as amended by the further plans and particulars submitted on the 4<sup>th</sup> September 2013, and 17<sup>th</sup> January 2014, except as may otherwise be required in order to comply with the following conditions.

Reason - In order to clarify the development to which this permission applies.

2. No development shall commence on site until a connection is made to the Limerick Main Drainage Scheme and is in operation to the Planning Authority's satisfaction as specified under planning permissions 06/1394 and 08/2320. The existing on site waste water system shall be decommissioned when the connection to the Limerick Main Drainage Scheme has been commissioned.

Reason - In the interest of orderly development.

3. a. No development shall commence on site until the developer(s) submits the following for the written agreement of the Planning Authority –
  - i. The developer engages the services of a suitably qualified person acceptable to the Planning Authority with professional indemnity insurance, who shall oversee all works on site as per the permission granted. Details shall be submitted for the written agreement of the Planning Authority.
  - ii. The developer shall notify the Planning Authority in writing at least one week prior to the commencement of any works to the site.
- b. On completion of works to the site a written report shall be submitted to the Planning Authority from the same suitably qualified person demonstrating that the development has been carried out in accordance with the planning permission granted.

The submission of such shall not absolve the developer of his responsibilities to construct and install infrastructural services in accordance with the requirements of this permission.

Reason – In the interest of orderly development.



4. Prior to commencement of development details of the proposed interceptor serving the "dirty area" shall be submitted for the written agreement of the Planning Authority. The interceptor must be in accordance with BS EN 858.

Reason - In the interest of proper planning and sustainable development.

5. Discharge from the truck/wheel wash shall be to the foul sewer via the proposed interceptor if detergents are not utilised in the washing process. If detergents are utilised, a zero-discharge recycling system shall be installed. No trucks other than those using the facility shall be permitted to use the wheel wash. Prior to commencement of development full details of the system shall be submitted for the written agreement of the Planning Authority.

Reason - In the interest of orderly development.

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**APPROPRIATE ASSESSMENT**

**STAGE 1 SCREENING**

**PROPOSED INCREASE IN WASTE ACCEPTANCE**

**MATERIALS RECOVERY FACILITY**

**GREENSTAR ENVIRONMENTAL SERVICES LTD**

**DOCK ROAD**

**LIMERICK**

**Prepared For: -**

Greenstar Environmental Services Ltd,  
Dock Road,  
Limerick.

**Prepared By: -**

O' Callaghan Moran & Associates,  
Granary House,  
Rutland Street,  
Cork.

**25<sup>th</sup> April 2013**

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

Project	Stage 1 Screening Assessment GES Limerick			
Client	GES			
Report No.	Date	Status	Prepared By	Reviewed By
12-4802203	25 <sup>th</sup> March	Draft	Jim O'Callaghan, MSc , CEnv	Micahel Watson, MA
	5 <sup>th</sup> April	Draft RevA		
	25 <sup>th</sup> April	Final		

For inspection purposes only.  
Consent of copyright owner required for any other use.

---

# TABLE OF CONTENTS

---

	<b><u>PAGE</u></b>
<b>1. INTRODUCTION .....</b>	<b>2</b>
1.1 <b>METHODOLOGY .....</b>	<b>3</b>
<b>2. DESCRIPTION OF PROJECT.....</b>	<b>4</b>
2.1 <b>SITE LOCATION .....</b>	<b>4</b>
2.2 <b>SITE LAYOUT.....</b>	<b>4</b>
2.3 <b>SITE OPERATIONS .....</b>	<b>5</b>
2.4 <b>DRAINAGE .....</b>	<b>5</b>
2.4.1 <i>Surface Water</i> .....	<b>5</b>
2.4.2 <i>Foul Water</i> .....	<b>8</b>
2.5 <b>PROPOSED DEVELOPMENT.....</b>	<b>9</b>
<b>3. NATURA 2000 SITES.....</b>	<b>10</b>
3.1 <b>NATURA 2000 SITES POTENTIALLY AFFECTED BY THE PROJECT.....</b>	<b>11</b>
3.2 <b>LOWER SHANNON SAC .....</b>	<b>12</b>
3.3 <b>RIVER SHANNON &amp; RIVER FERGUS SPA .....</b>	<b>13</b>
<b>4. LIKELY EFFECTS.....</b>	<b>15</b>
4.1 <b>PROPOSED DEVELOPMENT.....</b>	<b>15</b>
4.2 <b>DIRECT IMPACTS.....</b>	<b>15</b>
4.3 <b>INDIRECT IMPACTS.....</b>	<b>16</b>
4.4 <b>CUMULATIVE EFFECTS .....</b>	<b>16</b>
<b>5. SCREENING CONCLUSION &amp; STATEMENT.....</b>	<b>18</b>
 <b>APPENDIX 1</b> -            Limerick Docks Water Body Status Report	
 <b>APPENDIX 2</b> Site Synopses	
 <b>APPENDIX 3</b> Conservation Objectives	

---

# 1. INTRODUCTION

---

Greenstar Environmental Services Ltd (GES) intends to apply to for planning permission for its existing waste recovery and transfer facility in the townland of Ballykeefe, County Limerick. At a pre-application meeting with Limerick County Council, the Council requested that a Screening Assessment be prepared to inform the Appropriate Assessment of the application. GES appointed O'Callaghan Moran & Associates to carry out the assessment.

The European Union (EU) Habitats Directive (92/43/EC) and the EU Birds Directive (2009/147/EC) identify designated areas (Special Areas of Conservation (SAC) and Special Protection Areas (SPA) respectively) that are collectively known as Natura 2000 Sites. The Habitats Directive, which is implemented under the European Communities Birds and Natural Habitats) Regulations 2011 (S.I. No 477 of 2011), requires an “appropriate assessment” of the potential impacts any proposed development that may have an impact on the conservation objectives of any Natura 2000 site.

Article 6(3) of the Directive stipulates that *any plan or project not directly connected with or necessary to the management of a Natura 2000 site, but likely to have a significant effect thereon....shall be subject to appropriate assessment of its implications for the site in view of the site's conservation objectives.*

Guidance documents issued by Department of Environment, Heritage and Local Government and the National Parks and Wildlife Services recommend that the assessment be completed in a series of Stages, which comprise:

## *Stage 1: Screening*

The purpose of this Stage is to determine, on the basis of a preliminary assessment and objective criteria, whether a plan or project, alone and in combination with other plans or projects, could have significant effects on a Natura 2000 site in respect of the site's conservation objectives.

### *Stage 2: Appropriate Assessment*

This Stage is required if the Stage 1 Screening exercise identifies that the project is likely to have a significant impacts on a Natura 2000 site.

### *Stage 3 : Assessment of Alternative Solutions.*

If Stage 2 determines that the project will have an adverse impact upon the integrity of a Natura 2000 site, despite the implementation of mitigation measures, it must be objectively concluded that no alternative solutions exist before the plan can proceed.

### *Stage 4 : Compensatory Measures:*

Where no alternative solutions are feasible and where adverse impacts remain but imperative reasons of overriding public interest require the implementation of a project an assessment of compensatory measures that will effectively offset the damage to the Natura site 2000 is required.

## **1.1 Methodology**

The Screening Assessment was based on a site inspection and the proposed changes to facility operations. It followed the guidance presented in the “Assessment of Plans and Projects significantly affecting Natura 2000 sites, Methodological Guidance on the provisions of Articles 6(3) and 6(4) of the Habitats Directive 92/43/EEC” (2001); The DEHLG (2009, revised February 2010) Appropriate Assessment of Plans and Projects in Ireland and the NPWS (2010) Circular NPW 1/10 & PSSP 2/10 Appropriate Assessment under Article 6 of the Habitats Directive: Guidance for Planning Authorities.

---

## 2. DESCRIPTION OF PROJECT

---

### 2.1 Site Location

The GES facility is located in the townland of Ballykeefe, off the main N69 Limerick to Tralee road on Dock Road (Figure 2.1). It is in the northern end of an industrially zoned area and is bounded to the south, southeast and southwest by warehousing units, oil distribution centres and truck sales and repair and Cussen Crane Hire.

To the east and north is the Ballinacurra Creek, which is where the Ballynaclough River joins the Shannon. The lands north of the Ballinacurra and between it and the Shannon are undeveloped. The Limerick City Council wastewater treatment plant is to the west of the site and separated from it by an open field. Further west is Bunlickey Lake. .

### 2.2 Site Layout

The site layout is shown on Drawing No 002. The facility is accessed off the Dock Road by a private road common access road serving the facility and other occupiers of the industrial estate. The site encompasses 1.8ha, the vast majority of which is either paved or occupied by buildings.

There are two adjoining waste handling buildings (Building 1 and 2). Building 1 is currently used for sorting and compacting recyclables (paper, cardboard, plastics etc.) recovered from the incoming wastes. Building 2 is currently used for compacting and wrapping the mixed municipal solid wastes. There is a separate office building and adjoining vehicle and plant maintenance workshop near the site entrance. An electrical substation along the south-western boundary wall is owned by Electric Ireland.

The open yards are paved and are used for external waste storage bays (C&D, glass, metals, timber and baled waste), skip storage, truck parking and a vehicle wash area. There is palisade

security fence on the north, east and west boundaries, with block work walls along the south-western boundary south of Building 1 and west of the site offices and workshop.

## **2.3 Site Operations**

There are currently 20 full time employees based at the facility, including management, administration, general operatives and maintenance staff. The facility is authorised to operated seven days per week twenty four hours per day. At present, there are two eight hour shifts operating from 06:00 – 14:00 and 14:00 to 22:00.

The facility accepts and processes non hazardous mixed municipal solid waste and mixed and source segregated dry recyclables that are primarily collected in the Mid West Region.

The waste processing includes sorting of the mixed dry recyclables into separate categories (paper, plastic, cardboard), which are then compacted; the baling of the source segregated dry recyclables and the baling of the mixed municipal solid waste. The baled recyclables are sent to off-site recovery facilities for further processing, while the baled mixed municipal solid waste is sent to overseas waste to energy plants

## **2.4 Drainage**

### *2.4.1 Surface Water*

Surface water run-off is generated by rainfall on the roof of the offices and workshop building, the waste handling buildings and the paved open yard areas. The run-off from the paved yards is collected and directed through 2 No. three chamber oil interceptors before being discharged to a man made drain at the north-eastern site boundary. There is a shut off valve at the outlet from the last oil interceptor that can be closed in the event of an incident that has the potential to impact on surface water quality and contain the surface water within the site boundary.



Run-off from the main buildings discharges to manmade perimeter drain along the western boundary. The drainage layout is shown on Drawing No IE 580-002A.

The perimeter drains, which also take run-off from other occupants in the industrial estate, discharge to Bunlickey Lake. The water in the lake discharges to the Shannon River Estuary via valves and sluices that prevent tidal inflow.

The lower reaches of the Shannon are tidal and are part of the Shannon Transitional and Coastal Water Management Unit (WMU) designated in the Shannon River Basin District (ShIRBD) Management Plan prepared under the EU Water Framework Directive (WFD). The WMU comprises twenty Water Bodies and the stretch of the river to the north of the site is in the Limerick Dock Water Body.

Reports have been prepared on the 'Status' of each water body. Status means the condition of the water in a watercourse and is defined by its ecological status and chemical status, whichever is worse. Waters are ranked in one of five status classes, High, Good, Moderate, Poor and Bad. The WFD requires measures to ensure waters achieve at least 'Good Status' by specified period and that their current status does not deteriorate

The Limerick Dock Water Body Status Report, a copy of which is in Appendix 1, states that the water overall status is 'Good', with a High status for Biochemical Oxygen Demand, nutrients (phosphate and nitrogen) and dissolved oxygen. However, the overall chemical status is classified as 'Fail' and the water body is 'At Risk' of not achieving its restoration objective of reducing chemical pollution by 2021.

The risk assessment was prepared in 2008 and at that time the primary pressure on water quality identified in the Shannon Transitional and Coastal WMU Plan was combined sewer overflows and wastewater treatment plant overflows. Since then, the completion of the Limerick Main Drainage Scheme has significantly reduced the pressures on the Limerick Dock Water Body

The Waste Licence requires GES to monitor the quality of the surface water at specified locations monthly. These include the outlet from the interceptors (FE 1A) and in the receiving drain up (WS-9) and downstream (WS-10) of the discharge point. The locations

are shown on Figure 2.2 As the discharge is dependant on rainfall it is not always possible to collect samples at monthly intervals.

The monitoring parameters include pH, electrical conductivity, total suspended solids (TSS), ammonia, biochemical oxygen demand (BOD), Fats Oils and Grease (FOG), Mineral Oil, Total Organic Carbon (TOC) and dissolved metals (arsenic, cadmium, chromium, copper, lead, mercury, nickel and zinc).

The Waste Licence does not specify any emission limit values (ELVs) for the discharge, GES developed proposed ELVs for ammonia, BOD Mineral Oil and TSS, however these have not yet been agreed by the EPA. The monitoring results for 2012 are presented in Table 2.1

**Table 2.1 Water Quality Range 2012**

Parameter	Units	WS9	FE1A	WS10	ELV	EQS
pH	pH units	7.24-8.28	6.41-7.76	6.82-8.24	-	
BOD	mg/l	1-7	35-176	2-89	25	1.5
TSS	mg/l	1-12	50-130	2-51	60	
Ammonia	mg/l	0.27-1	0.14-3.64	0.04-<1	4	0.065
FOG	mg/l	<0.01-<1	0.29-17.2	<0.01,3.3	-	-
Mineral Oils	mg/l	<0.01-<1	<0.01-2.03	<0.01-<7	5	
TOC	mg/l	3.57-18	22.63-48	4.25-20	-	-
Arsenic	ug/l	0.001-3	0.002-5	0.001<3		25
Cadmium	ug/l	<0.03-0.5	>0.03-0.02	0-<1	-	5
Chromium	ug/l	<1.5-2	0.5-2.4	0.2-<1.5	-	30
Copper	ug/l	<0.2-13	<0.2-16	<0.2-13	-	30
Mercury	ug/l	<0.0001-1	<0.0001-<1	<0.0001-<1	-	1
Nickel	ug/l	<0.2-2	<0.2-14.1	<0.2-2.4	-	20
Lead	ug/l	0.2-<2	0.2-<5	<0.2-<5	-	10
Zinc	ug/l	<0.2-11	<0.2-47.5	<0.2-15	-	100

For those parameters for which ELVs have not been established the Environmental Quality Standards (EQS) specified for 'Good Status' in the Environmental Objectives (Surface Water) Regulations 2009 (S.I. No.272 of 2009) are provided. The EQS are not emission

limit values, but are the concentrations that must be achieved in a water body, taking into consideration the available assimilative capacity, if the water body is to meet the objectives set for the water body.

The monitoring indicates that, with the exception of BOD and TSS, all of the parameters are below the proposed ELV and significantly below the EQS. In particular mineral oils have never been detected. It is noted that the BOD and ammonia levels in the drain upstream of the discharge point exceed the EQS.

In 2012, GES conducted an extensive CCTV survey of the surface water drainage system. The survey identified a number of defects in the surface water lines, some small cracks in the first chamber of the interceptor and further cracks in the pipeline connecting the final chamber of the interceptors to the discharge point. These defects were repaired in May 2012.

#### *2.4.2 Foul Water*

Sanitary wastewater and wastewater from the vehicle wash area is treated in to the on-site Klargester Biodisc wastewater treatment plant. The vehicle wastewater passes through a grit trap and oil interceptor before connecting to the Klargester. Sanitary wastewater from the neighbouring Cussen Crane Hire Yard is also connected to the Klargester. The treated effluent discharges to a percolation area and the quality of the discharge is monitored in accordance with the requirements of the Waste Licence.

In 2012 GES commissioned a detailed assessment of the operation of the treatment plant. The assessment established that the average daily discharge to the percolation area is 0.4m<sup>3</sup>/day. Taking into consideration rainfall on the percolation area, the total hydraulic loading is 0.483m<sup>3</sup>/day. The effluent quality monitoring has established that the quality meets the recommended minimum performance standards set by the EPA and are within the manufacturer's design standards.

It is a condition of the waste licence that discharge foul water and sewage from the site must be to the Council's foul sewer, following the completion of the Limerick Main Drainage Scheme, subject to the approval of the Sanitary Authority-Limerick City Council.


In 2009, the City Council gave its approval in principle to the connection to the municipal wastewater treatment plant, however due to difficulties in obtaining way leaves to install the sewer line, the connection could not be completed at that time. GES is currently engaged with both the City and County Councils regarding the connection and the necessary wayleaves and it is expected that the connection will be completed by ??? Following this the on-site wastewater treatment plant will be decommissioned.

## **2.5 Proposed Development**

GES intends to increase the amount of waste that can be accepted to 130,000 tonnes/year. The proposed increase is to allow GES compete for business in domestic and commercial waste collection market and offer waste treatment services to authorised waste collectors in the Mid West and adjoining Regions.

There will be no change to either the types of waste accepted, or the way the waste is handled, processed and stored. The only change will be an increase in the number of vehicles that bring the unprocessed waste to the site and remove the processed materials.

For inspection purposes only.  
Consent of copyright owner required for any other use.



For inspection purposes only. Not for any other use.  
 Consent of copyright owner required.


**O' Callaghan Moran & Associates.**  
 Granary House, Rutland Street,  
 Cork, Ireland.  
 Tel. (021) 4321521 Fax. (021) 4321522  
 email : info@ocalaghanmoran.com

CLIENT  
**Greenstar Environmental Services Ltd**

details

Figure No.  
**2.1**

This drawing is the property of O'Callaghan Moran & Associates and shall not be used, reproduced or disclosed to anyone without the prior written permission of O'Callaghan Moran & Associates and shall be returned upon request.


TITLE  
**WASTE LICENCE AREA**

SCALE	REV.
1:750 A4	A



NOTES

- Monitoring locations  
Grid references  
Eastings Northings
- DM-1 - 154913, 155617
  - DM-2 - 154842, 155714
  - DM-3 - 155003, 155750
  - Ni-1 - 154840, 155712
  - Ni-2 - 154920, 155745
  - Ni-3 - 154910, 155618
  - FE-1A - 154974, 155769
  - WS-9 - 155003, 155750
  - WS-10 - 154922, 155776
  - FE-2A - 154858, 155720
  - GW-M1 - 154923, 155667
  - GW-M2 - 154920, 155770
  - GW-M3 - 155049, 155684



- SITE BOUNDARY (RED)
- SURFACEWATER DRAINAGE
- FOULWATER DRAINAGE
- NOISE MONITORING LOCATION
- ⊕ DUST MONITORING LOCATION
- ⊠ GROUNDWATER MONITORING LOCATION
- ⊕ FOULWATER MONITORING LOCATION

For inspection purposes only.  
Consent of copyright owner required for any other use.

A	02/05/13	PRELIMINARY ISSUE	MW	JOC	∞
REV	DATE	DESCRIPTION	DRN	CHKD	APP


**OCA**  
O'Callaghan Moran & Associates.  
Granary House, Rutland Street,  
Cork, Ireland.  
Tel. (021) 4321521 Fax (021) 4321522  
email : oam@indigo.ie

This drawing is the property of O'Callaghan Moran & Associates and shall not be used, reproduced or disclosed in any way without the prior written permission of O'Callaghan Moran & Associates and shall be returned upon request.


CLIENT  
**GES Ltd**

TITLE  
**GES Limerick  
Monitoring Locations  
W0082-02**

SCALE	DRAWING No.	REV.
NTS A3	2.2	A



— SITE BOUNDARY (RED)  
- - - SURFACEWATER DRAINAGE  
- - - FOUWATER DRAINAGE



REV	DATE	DESCRIPTION	DRN	CHKD	APP
A	01/05/13	PRELIMINARY ISSUE	MW	JOC	**

This drawing is the property of O'Callaghan Moran & Associates and shall not be used for any other project without the written permission of O'Callaghan Moran & Associates and shall be returned upon request.  
 O'Callaghan Moran & Associates  
 Gentry House, Rutland Street,  
 Cork, Ireland.  
 Tel: (021) 4321521 Fax: (021) 4321522  
 email: info@ocallaghantomoran.com

**GES Limerick**  
 CLIENT

**SITE LAYOUT & DRAINAGE**  
 DRAWING No. 002  
 SCALE 1:500 A

---

### 3. NATURA 2000 SITES

---

SACs are selected for the conservation and protection of habitats listed on Annex I and species (other than birds) listed on Annex II of the Habitats Directive, and their habitats. The habitats on Annex I require special conservation measures. SPAs are selected for the conservation and protection of bird species listed on Annex I of the Birds Directive and regularly occurring migratory species, and their habitats, particularly wetlands. The selected habitats and species are termed Qualifying Interests

A statement of Conservation Objectives is prepared for each designated site which identifies the qualifying interests or conservation features. The Conservation Objectives are intended to ensure that the relevant habitats and species present on a site are maintained, and where necessary restored, at a Favourable Conservation Status.

Favourable Conservation Status of a habitat, as defined in 2011 Birds and Natural Habitats Regulations, is when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable

Conservation Status of a species is when:

- the Favourable population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats,
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

A list of designated Natura 2000 sites within 15 km of the facility is given in Table 3.1



**Table 3.1.** Natura 2000 Sites Within 15 km of the AES Facility

Site	Code	Distance
<b>SAC</b>		
Lower River Shannon	002165	400m to the north of the site.
Glenorma Wood	1013	11.4 km to the north east
Ratty River Cave	2316	14.5 km to the north east
Danes Hole Poulnalecka	0030	15 km to the north
Tory Hill	0439	13 km to the south
Askeaton Fen Complex	002279	14km to the south west
<b>SPA</b>		
River Shannon & River Fergus	004077	400 m to the north of the site

### 3.1 Natura 2000 Sites Potentially Affected by the Project

The facility is not located in or immediately adjacent to a Natura 2000 Site. The closest Natura 2000 Sites are the Lower River Shannon SAC and the River Shannon & River Fergus SPA, which are 400m to the north.

Stormwater run-off from the site discharges to Bunlickey Lake which is in the River Shannon & River Fergus SPA and is hydraulically connected to the River Shannon. The remaining Sites are between 10 and 15 km from the facility and there is no pathway by which the current and proposed site activities can impact on these Sites

### 3.2 Lower Shannon SAC

The Site Synopsis for the Lower Shannon SAC that lists the full Qualifying Interests are in Appendix 2, and the Conservation Objective are in Appendix 3 and the information is summarised below.

#### *Qualifying Interests*

The Lower Shannon SAC is selected for the following habitats listed in Annex 1 of the Habitats Directive: lagoons and alluvial wet woodlands, floating river vegetation, *Molinia* meadows, estuaries, tidal mudflats, Atlantic salt meadows, Mediterranean salt meadows, *Salicornia* mudflats, sand banks, perennial vegetation of stony banks, sea cliffs, reefs and large shallow inlets and bays all habitats

The site is also selected for the following species listed in Annex II of the Directive – Bottle Nosed Dolphin, Sea Lamprey, River Lamprey, Brook Lamprey, Freshwater Pearl Mussel, Atlantic Salmon and Otter.

#### *Conservation Objectives*

The conservation objectives are to maintain or restore the favorable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected??

- [1029] Freshwater pearl mussel *Margaritifera margaritifera*
- [1095] sea lamprey *Petromyzon marinus*
- [1096] Brook Lamprey *Lampetra planeri*
- [1099] River Lamprey *Lampetra fluviatilis*
- [1106] Atlantic Salmon *Salmo salar* (only in fresh water)
- [1110] sandbanks which are slightly covered by sea water all the time
- [1130] Estuaries
- [1140] Mudflats and sandflats not covered by seawater at low tide
- [1150] \*Coastal lagoons
- [1160] Large shallow inlets and bays

- [1170] Reefs
- [1220] Perennial vegetation of stony banks
- [1230] vegetated sea cliffs of the atlantic and Baltic coasts
- [1310] *Salicornia* and other annuals colonizing mud and sand
- [1330] Atlantic salt meadows (*glauco-puccinellietalia maritimae*)
- [1349] Bottlenose Dolphin *Tursiops truncatus*
- [1355] Otter *lutra lutra*
- [1410] Mediterranean salt meadows (*Juncetalia maritima*)
- [3260] water courses of plain to montane levels with the *ranunculion fluitantis* and *Callitricho- Batrachion* vegetation
- [91EO] \*Alluvial forests with *Alnus gutinosa* and *Fraxinus excelsior* (*Alno-padion*, *alnion incanae*, *salicion albae*)

### 3.3 River Shannon & River Fergus SPA

The Site Synopsis and for the River Shannon & River Fergus SPA listing the Qualifying Interests and the Conservation Objective are in Appendix 2 and are summarised below.

#### *Qualifying Interests*

The Shannon and Fergus Estuaries SPA comprises the entire estuarine habitat west of Limerick City and south of Ennis extending approximately 25 km west to Killadysert and Foynes on the north and south shores of the Shannon.

The Site is the most important coastal wetland site in the country and regularly supports in excess of 50,000 wintering waterfowl. Other species occurring include Common Cockle (*Cerastoderma edule*), Lugworm (*Arenicola marina*), polychaete *Nephtys hombergii*, gastropod *Hydrobia ulvae* and the crustacean *Corophium volutator*. Eelgrass (*Zostera* spp.) is present in places, along with green algae (e.g. *Ulva* spp. and *Enteromorpha* spp.). The Site also has extensive intertidal flats, which is a listed habitat in Annex 1 of the Habitats Directive.

## Conservation Objectives

The conservation objectives are to maintain or restore the favorable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SPA has been selected:

- [A017] Cormorant *phalacrocorax carba*
- [A038] Whooper swan *Cygnus Cygnus*
- [A046] Light-bellied Brent Goose *Branta bernicla hrota*
- [A048] Shelduck *Tadorna tadorna*
- [A050] Wigeon *Anas Penelope*
- [A052] Teal *Anas crecca*
- [A054] Pintail *Anas acuta*
- [A056] Shoveler *Anas clypeata*
- [A062] Scaup *Aythya marila*
- [A137] Ringed Plover *chahrius hiaticula*
- [A140] golden plover *pluvialis apricaria*
- [A141] Grey plover *pluvialis apricaria*
- [A142] Lapwing *Vanellus vanellus*
- [A143] Knot *Calidris canuts*
- [A149] Dunlin *Calidris alpine*
- [A156] Blacktailed Godwit *Limosa limosa*
- [A157] Bar-tailed godwit *limosa lapponica*
- [A160] Curlew *Numenius arquata*
- [A162] Redshank *Tringa tetanus*
- [A164] Greenshank *Trina nebularia*
- [A179] Black headed Gull *Chroicocephalus ridibundus*
- [A999] Wetlands

---

## 4. LIKELY EFFECTS

---

### 4.1 Proposed Development

The proposed increase in the annual waste throughput will not require the expansion of the site, the construction/provision of any new buildings/structures, or any alteration to the existing site layout and operations.

There will be no change to the waste acceptance and operational hours and it will not require the use of any new raw materials that have the potential to cause contamination. It will not result in any new or additional abstraction from groundwater or surface water. It will not give rise to any new emissions to surface water or sewer, nor will it contribute to increased noise, dust and odour emissions or illumination.

### 4.2 Direct Impacts

The GES facility is not located within any designated Natura 2000 Site and therefore the proposed changes will not result in any direct habitat loss or fragmentation of either the Lower River Shannon SAC or the River Shannon and River Fergus Estuaries SPA.

These Natura 2000 Sites are approximately 400m to the north and west of the GES facility. The facility is extensively developed and almost entirely covered with paving and buildings, which means it does not support the species for which the Natura 2000 sites were selected.

Based on the above, the project does not present any risk of a direct adverse affect on either the habitats or species for which the Natura 2000 Sites were selected.

### **4.3 Indirect Impacts**

There is the potential for indirect impacts on the Natura 2000 Sites, as surface water run-off from the yards and roofs discharges to the Bunlickey Lake, which is part of the River Shannon & Fergus SPA and hydraulically connected to the River Shannon via sluices. However, the project will not result in any changes to either the volume or quality of the surface water run-off from the facility and therefore will have no impact of the Natural 2000 Sites.

Disturbance impacts are considered with regard to the potential for effects on the Annex II species for which the Lower River Shannon SAC is designated and the bird species listed as special conservation interests of the River Shannon and River Fergus Estuaries SPA.

The GES facility is located within an industrial estate and is 2km west of Limerick Docks. There are extensive and ongoing traffic movements, artificial lighting and noise emissions associated with both areas. It must be noted that the presence of the listed species of conservation interest within the environs of Limerick City indicates they have become acclimatised to the background levels of disturbance.

The project does not require the provision of any new plant and equipment or changes to the operational hours therefore there will be no additional sources of disturbance to the listed species present in both the SAC and SPA.

### **4.4 Cumulative Effects**

Recent projects completed within the SAC include the River Fergus Lower (Ennis) Drainage Scheme and maintenance works carried out by the OPW on upstream of Limerick City and on the River Maigue at Adare in 2010. Maintenance works are being undertaken in the Abbey River corridor which will include dredging from the Park Canal confluence to the confluence with the Shannon.

Point and diffuse sources of water pollution in the urban area comprise a cumulative pressure on the conservation interests of the SAC, where Annex II aquatic species are considered to be under stress due to poor background water quality. The proposed increase in the amount of waste accepted will not result in any changes to either the volume or quality of the surface water run-off

that therefore and will not contribute to any significant cumulative impact on the Natura 2000 Sites

The proposed changes does not involve the construction of new buildings, the introduction of new plant an equipment or the changes to the operational hours, and therefore will not add to the cumulative disturbance effects on the Natura 2000 Sites.

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

---

## 5. SCREENING CONCLUSION & STATEMENT

---


The proposed increase in the annual waste throughput will not result in any new or additional emissions/disturbance that could present a significant risk to the Qualifying Interests and Conservation Objectives of either the Lower Shannon SAC or the Shannon and Fergus Estuaries SPA. Therefore a Natura Impact Statement is not required.

*For inspection purposes only.  
Consent of copyright owner required for any other use.*




# APPENDIX 1

For inspection purposes only.  
Consent of copyright owner required for any other use.



## Full Report for Waterbody Limerick Dock



River Basin Management Plans (RBMPs) have been published for all River Basin Districts in Ireland in accordance with the requirements of the Water Framework Directive. The WaterMaps viewer is an integral part of the River Basin Management Plan and provides access to information at individual waterbody level and at Water Management Unit level for all the River Basin Districts in Ireland.

The following report provides summary plan information about the selected waterbody (indicated by the pin in the map above) relating to its status, risks, objectives, and measures proposed to retain status where this is adequate, or improve it where necessary. Waterbodies can relate to surface waters (these include rivers, lakes, estuaries [transitional waters], and coastal waters), or to groundwaters. Other relevant information not included in this report can be viewed using the WaterMaps viewer, including areas listed in the Register of Protected Areas.

You will find brief notes at the bottom of some of the individual report sheets that will help you in interpreting the information presented. More detailed information can be obtained in relation to all aspects of the RBMPs at [www.wfdireland.ie](http://www.wfdireland.ie).

Date Reported to Europe: July 2010

Date Report Created 11/07/2012



**Summary Information:**

**Water Management Unit:** N/A  
**WaterBody Category:** Transitional Waterbody  
**WaterBody Name:** Limerick Dock  
**WaterBody Code:** IE\_SH\_060\_0900  
**Overall Status:** Good  
**Overall Objective:** Restore 2021  
**Overall Risk:** 1a At Risk  
**Heavily Modified:** Yes



Report data based upon final RBMP, 2009-2015.

The information provided above is a summary of the principal findings related to the selected waterbody. Further details and explanation of individual elements of the report are outlined in the following pages.

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

Date Reported to Europe: July 2010

Date Report Created 11/07/2012



**Status Report**

**Water Management Unit:** N/A  
**WaterBody Category:** Transitional Waterbody  
**WaterBody Name:** Limerick Dock  
**WaterBody Code:** IE\_SH\_060\_0900  
**Overall Status Result:** Good  
**Heavily Modified:** Yes



<b>Status Element Description</b>		<b>Result</b>
<b>Status information</b>		
DIN	Dissolved Inorganic Nitrogen status	<span style="background-color: green; color: white; padding: 2px;">Good</span>
MRP	Molybdate Reactive Phosphorus status	<span style="background-color: #333399; color: white; padding: 2px;">High</span>
DO	Dissolved oxygen as per cent saturation status	<span style="background-color: #333399; color: white; padding: 2px;">High</span>
BOD	Biochemical Oxygen Demand (5-days) status	<span style="background-color: #333399; color: white; padding: 2px;">High</span>
PHY	Macroalgae - phytobiomass status	<span style="background-color: #333399; color: white; padding: 2px;">High</span>
OPP	Macroalgae - opportunistic algae status	N/A
RSL	Macroalgae - reduced species list status	N/A
ANG	Angiosperms - Seagrass and Saltmarsh status	N/A
BIN	Benthic Invertebrates status	N/A
FIS	Fish status	<span style="background-color: green; color: white; padding: 2px;">Good</span>
HYD	Hydrology status	N/A
MOR	Morphology status	Less than Good (pHMWB)
SP	Specific Pollutant Status	Pass
PAS	Overall protected area status	At least good
ES	Ecological Status	<span style="background-color: green; color: white; padding: 2px;">Good</span>
CS	Chemical Status	Fail
SWS	Surface Water Status	N/A
EXT	Extrapolated status	N/A
DON	Donor water bodies	N/A

For inspection purposes only.  
Consent of copyright owner required for any other use.

Date Reported to Europe: July 2010

Date Report Created 11/07/2012



n/a - not assessed

**Status**

By 'Status' we mean the condition of the water in the waterbody. It is defined by its chemical status and its ecological status, whichever is worse. Waters are ranked in one of 5 status classes: High, Good, Moderate, Poor, Bad. However, not all waterbodies have been monitored, and in such cases the status of a similar nearby waterbody has been used (extrapolated) to assign status. If this has been done the first line of the status report shows the code of the waterbody used to extrapolate.

You can read more about status and how it is measured in our RBMP Document Library at [www.wfdireland.ie](http://www.wfdireland.ie) (Directory 15 Status).

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

Date Reported to Europe: July 2010

Date Report Created 11/07/2012




**Risk Report**

**Water Management Unit:** N/A  
**WaterBody Category:** Transitional Waterbody  
**WaterBody Name:** Limerick Dock  
**WaterBody Code:** IE\_SH\_060\_0900  
**Overall Risk Result:** 1a At Risk  
**Heavily Modified:** Yes



<b>Risk Test Description</b>		<b>Risk</b>
<b>Hydrology</b>		
THY1	Water balance - Abstraction	<span style="background-color: purple; color: white; padding: 2px;">2b</span> Not At Risk
<b>Marine Direct Impacts</b>		
TMDI 1	Dangerous Substances	N/A
TMDI 2	OSPAR	N/A
TMDI 3	UWWT Regs Designations	N/A
TMDI 0	Marine Direct Impacts Overall - Worst Case	N/A
<b>Morphological Risk Sources</b>		
TM1	Channelisation	N/A
TM2	Deposition	N/A
TM3	Coastal Defences	N/A
TM4	Impoundments	N/A
TM5a	Built Structures - Port Tonnage	N/A
TM5b	Built Structures - Industrial Intakes	N/A
TM6	Intensive Landuse	N/A
TMO	Morphology Overall - Worst Case	N/A
TMO	Overall (MIMAS) Morphological Risk - Worst Case (2008)	N/A
<b>Overall Risk</b>		
RA	Transitional Overall - Worst CaseOverall (MIMAS) Morphological Risk - Worst Case (2008)	<span style="background-color: purple; color: white; padding: 2px;">1a</span> At Risk
<b>Point / MDI Worst Case</b>		
TPOL	Worst case of Point Overall and MDI OverallOverall (MIMAS) Morphological Risk - Worst Case (2008)	<span style="background-color: purple; color: white; padding: 2px;">1a</span> At Risk

Consent of copyright owner required for any other use.



Point Risk Sources		
TP1	WWTPs (2008)	2b Not At Risk
TP2	CSOs	1a At Risk
TP3	IPPCs (2008)	2b Not At Risk
TP4	Section 4s (2008)	2b Not At Risk
TP5	WTPs/Mines/Quarries/Landfills	N/A
TPO	Overall Risk from Point Sources - Worst Case (2008)	1a At Risk

**Risk**

By 'risk' we mean the risk that a waterbody will not achieve good ecological or good chemical status/potential at least by 2015. To examine risk the various pressures acting on the waterbody were identified along with any evidence of impact on water status. Depending on the extent of the pressure and its potential for impact, and the amount of information available, the risk to the water body was placed in one of four categories: 1a at risk; 1b probably at risk; 2a probably not at risk; 2b not at risk. Note that '2008' after the risk category means that the risk assessment was revised in 2008. All other risks were determined as part of an earlier risk assessment in 2005.

You can read more about risk assessment in our 'WFD Risk Assessment Update' document in the RBMP document library, and other documents at [www.wfdireland.ie](http://www.wfdireland.ie) (Directory 31 Risk Assessments).

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

Date Reported to Europe: July 2010

Date Report Created 11/07/2012



**Objectives Report**

**Water Management Unit:** N/A  
**WaterBody Category:** Transitional Waterbody  
**WaterBody Name:** Limerick Dock  
**WaterBody Code:** IE\_SH\_060\_0900  
**Overall Objective:** Restore 2021  
**Heavily Modified:** Yes



<b>Objectives Description</b>		<b>Result</b>
<b>Extended timescale information</b>		
E1	Extended timescales due to time requirements to upgrade WWTP discharges	No Status
E2	Extended timescales due to delayed recovery of chemical pollution and chemical status failures	2021
E3	Extended timescales due to winter dissolved nitrogen exceedances	No Status
E4	Extended timescales due to time requirements for status recovery	No Status
E5	Extended timescales from Northern Ireland Environment Agency	No Status
E0V	Overall extended timescale - combination of all extended timescales fields	2021
<b>Objectives information</b>		
OB1	Prevent deterioration objective	No Status
OB2	Restore at least good status objective	No Status
OB3	Reduce chemical pollution objective	Restore 2021
OB4	Protected areas objective	Protect
OBO	Overall objectives	Restore 2021

**Extended timescales**

Extended timescales have been set for certain waters due to technical, economic, environmental or recovery constraints. Extended timescales are usually of one planning cycle (6 years, to 2021) but in some cases are two planning cycles (to 2027).

**Objectives**

In general, we are required to ensure that our waters achieve at least good status/potential by 2015, and that their status does not deteriorate. Having identified the status of waters (this is given earlier in this report), the next stage is to set objectives for waters. Objectives consider waters that require protection from deterioration as well as waters that require restoration and the timescales needed for recovery. Four default objectives have been set initially:-

- Prevent Deterioration*
- Restore Good Status*
- Reduce Chemical Pollution*
- Achieve Protected Areas Objectives*

These objectives have been refined based on the measures available to achieve them, the latter's likely effectiveness, and consideration of cost-effective combinations of measures. Where it is considered necessary extended deadlines have been set for achieving objectives in 2021 or 2027.

Date Reported to Europe: July 2010

Date Report Created 11/07/2012





**Measures Report**

**Water Management Unit:** N/A  
**WaterBody Category:** Transitional Waterbody  
**WaterBody Name:** Limerick Dock  
**WaterBody Code:** IE\_SH\_060\_0900  
**Heavily Modified:** Yes



	<b>Measures Description</b>	<b>Applicable</b>
BC	Total number of basic measures which apply to this waterbody	16
BW	Directive - Bathing Waters Directive	No
BIR	Directive - Birds Directive	Yes
HAB	Directive - Habitats Directive	Yes
MAE	Directive - Major Accidents and Emergencies Directive	Yes
EIA	Directive - Environmental Impact Assessment Directive	Yes
UWT	Directive - Urban Waste Water Treatment Directive	No
PPP	Directive - Plant Protection Products Directive	Yes
NIT	Directive - Nitrates Directive	Yes
IPC	Directive - Integrated Pollution Prevention Control Directive	Yes
POI	Other Stipulated Measure - Control of point source discharges	Yes
DIF	Other Stipulated Measure - Control of diffuse source discharges	Yes
PS	Other Stipulated Measure - Control of priority substances	Yes
MOD	Other Stipulated Measure - Controls on physical modifications to surface waters	Yes
OA	Other Stipulated Measure - Controls on other activities impacting on water status	Yes
AP	Other Stipulated Measure - Prevention or reduction of the impact of accidental pollution incidents	Yes
TP1	WSIP - Agglomerations with treatment plants requiring capital works	No
TP2	WSIP - Agglomerations with treatment plants requiring further investigation prior to capital works	No
TP3	WSIP - Agglomerations requiring the implementation of actions identified in Shellfish PRPs	No
TP4	WSIP - Agglomerations with treatment plants requiring improved operational performance	No
TP5	WSIP - Agglomerations requiring investigation of CSOs	No
TP6	WSIP - Agglomerations where existing treatment capacity is currently adequate but predicted loadings would result in overloading	No
OTS	On-site waste water treatment systems	Yes
SHE	Shellfish Pollution Reduction Plan	No
IPR	IPPC licences requiring review	Yes
WPR	Water Pollution Act licences requiring review	Yes

Date Reported to Europe: July 2010

Date Report Created 11/07/2012



HQW	Protect high quality waters	No
-----	-----------------------------	----

**Measures**

Measures are necessary to ensure that we meet the objectives set out in the previous page of this report. Many measures are already provided for in national legislation and must be implemented. Other measures have been recently introduced or are under preparation. A range of additional potential measures are also being considered but require further development. Any agreed additional measures can be introduced through the update of Water Management Unit Action Plans during the implementation process.

You can read more about Basic Measures in 'River Basin Planning Guidance' and in other documents in our RBMP Document Library at [www.wfdireland.ie](http://www.wfdireland.ie).

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

Date Reported to Europe: July 2010

Date Report Created 11/07/2012

# **APPENDIX 2**

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

## SITE SYNOPSIS

**SITE NAME : LOWER RIVER SHANNON**

**SITE CODE : 002165**

This very large site stretches along the Shannon valley from Killaloe to Loop Head/ Kerry Head, a distance of some 120 km. The site thus encompasses the Shannon, Feale, Mulkear and Fergus Estuaries, the freshwater lower reaches of the River Shannon (between Killaloe and Limerick), the freshwater stretches of much of the Feale and Mulkear catchments and the marine area between Loop Head and Kerry Head. The Shannon and Fergus flow through Carboniferous limestone as far as Foynes, but west of Foynes Namurian shales and flagstones predominate (except at Kerry Head, which is formed from Old Red Sandstone). The eastern sections of the Feale catchment flow through Namurian Rocks and the western stretches through Carboniferous Limestone. The Mulkear flows through Lower Palaeozoic Rocks in the upper reaches before passing through Namurian Rocks, followed by Lower Carboniferous Shales and Carboniferous Limestone. The Mulkear River itself, immediately north of Pallas Green, passes through an area of Rhyolites, Tuffs and Agglomerates. Rivers within the sub-catchment of the Feale include the Galey, Smearlagh, Oclagh, Allaughaun, Owveg, Clydagh, Caher, Breanagh and Glenacarne. Rivers within the sub-catchment of the Mulkear include the Killeenagarraiff, Annagh, Newport, the Dead River, the Bilboa, Glashacloonaraveela, Gortnageragh and Cahernahallia.

The site is a candidate SAC selected for lagoons and alluvial wet woodlands, both habitats listed on Annex I of the E.U. Habitats Directive. The site is also selected for floating river vegetation, *Molinia* meadows, estuaries, tidal mudflats, Atlantic salt meadows, Mediterranean salt meadows, *Salicornia* mudflats, sand banks, perennial vegetation of stony banks, sea cliffs, reefs and large shallow inlets and bays all habitats listed on Annex I of the E.U. Habitats Directive. The site is also selected for the following species listed on Annex II of the same directive – Bottle-nosed Dolphin, Sea Lamprey, River Lamprey, Brook Lamprey, Freshwater Pearl Mussel, Atlantic Salmon and Otter.

The Shannon and Fergus Estuaries form the largest estuarine complex in Ireland. They form a unit stretching from the upper tidal limits of the Shannon and Fergus Rivers to the mouth of the Shannon estuary (considered to be a line across the narrow strait between Kilcredaun Point and Kilconly Point). Within this main unit there are several tributaries with their own 'sub-estuaries' e.g. the Deel River, Mulkear River, and Maigue River. To the west of Foynes, a number of small estuaries form indentations in the predominantly hard coastline, namely Poulnasherry Bay, Ballylongford Bay, Clonderalaw Bay and the Feale or Cashen River Estuary.

Both the Fergus and inner Shannon estuaries feature vast expanses of intertidal mudflats, often fringed with saltmarsh vegetation. The smaller estuaries also feature mudflats, but have their own unique characteristics, e.g. Poulnasherry Bay is stony and unusually rich in species and biotopes. Plant species are typically scarce on the mudflats, although there are some Eel-grass beds (*Zostera* spp.) and patches of green

algae (e.g. *Ulva* sp. and *Enteromorpha* sp.). The main macro-invertebrate community, which has been noted from the inner Shannon and Fergus estuaries, is a *Macoma-Scrobicularia-Nereis* community.

In the transition zone between mudflats and saltmarsh, specialised colonisers of mud predominate: swards of Common Cord-grass (*Spartina anglica*) frequently occur in the upper parts of the estuaries. Less common are swards of Glasswort (*Salicornia europaea* agg.). In the innermost parts of the estuaries, the tidal channels or creeks are fringed with species such as Common Reed (*Phragmites australis*) and Club-rushes (*Scirpus maritimus*, *S. tabernaemontani* and *S. triquetrus*). In addition to the nationally rare Triangular Club-rush (*Scirpus triquetrus*), two scarce species are found in some of these creeks (e.g. Ballinacurra Creek): Lesser Bulrush (*Typha angustifolia*) and Summer Snowflake (*Leucojum aestivum*).

Saltmarsh vegetation frequently fringes the mudflats. Over twenty areas of estuarine saltmarsh have been identified within the site, the most important of which are around the Fergus Estuary and at Ringmoylan Quay. The dominant type of saltmarsh present is Atlantic salt meadow occurring over mud. Characteristic species occurring include Common Saltmarsh Grass (*Puccinellia maritima*), Sea Aster (*Aster tripolium*), Thrift (*Armeria maritima*), Sea-milkwort (*Glaux maritima*), Sea Plantain (*Plantago maritima*), Red Fescue (*Festuca rubra*), Creeping Bent (*Agrostis stolonifera*), Saltmarsh Rush (*Juncus gerardi*), Long-bracted Sedge (*Carex extensa*), Lesser Sea-spurrey (*Spergularia marina*) and Sea Arrowgrass (*Triglochin maritima*). Areas of Mediterranean salt meadows, characterised by clumps of Sea Rush (*Juncus maritimus*) occur occasionally. Two scarce species are found on saltmarshes in the vicinity of the Fergus Estuary: a type of robust Saltmarsh-grass (*Puccinellia foucaudii*), sometimes placed within the compass of Common Saltmarsh-grass (*Puccinellia maritima*) and Hard-grass (*Parapholis strigosa*).

Saltmarsh vegetation also occurs around a number of lagoons within the site. The two which have been surveyed as part of a National Inventory of Lagoons are Shannon Airport Lagoon and Clooncneen Pool. Clooncneen Pool (4-5 ha) is a natural sedimentary lagoon impounded by a low cobble barrier. Seawater enters by percolation through the barrier and by overwash. This lagoon represents a type which may be unique to Ireland since the substrate is composed almost entirely of peat. The adjacent shore features one of the best examples of a drowned forest in Ireland. Aquatic vegetation in the lagoon includes typical species such as Beaked Tasselweed (*Ruppia maritima*) and green algae (*Cladophora* sp.). The fauna is not diverse, but is typical of a high salinity lagoon and includes six lagoon specialists (*Hydrobia ventrosa*, *Cerastoderma glaucum*, *Lekanesphaera hookeri*, *Palaemonetes varians*, *Sigara stagnalis* and *Enochrus bicolor*). In contrast, Shannon Airport Lagoon (2 ha) is an artificial saline lake with an artificial barrier and sluiced outlet. However, it supports two Red Data Book species of Stonewort (*Chara canescens* and *Chara cf. connivens*).

Most of the site west of Kilcredaun Point/Kilconly Point is bounded by high rocky sea cliffs. The cliffs in the outer part of the site are sparsely vegetated with lichens, Red Fescue, Sea Beet (*Beta vulgaris*), Sea Campion (*Silene maritima*), Thrift and Plantains (*Plantago* spp.). A rare endemic Sea Lavender (*Limonium recurvum* subsp.

*pseudotranswallinum*) occurs on cliffs near Loop Head. Cliff-top vegetation usually consists of either grassland or maritime heath. The boulder clay cliffs further up the estuary tend to be more densely vegetated, with swards of Red Fescue and species such as Kidney Vetch (*Anthyllis vulneraria*) and Bird's-foot Trefoil (*Lotus corniculatus*).

The site supports an excellent example of a large shallow inlet and bay. Littoral sediment communities in the mouth of the Shannon Estuary occur in areas that are exposed to wave action and also in areas extremely sheltered from wave action. Characteristically, exposed sediment communities are composed of coarse sand and have a sparse fauna. Species richness increases as conditions become more sheltered. All shores in the site have a zone of sand hoppers at the top and below this each of the shores has different characteristic species giving a range of different shore types in the pcSAC.

The intertidal reefs in the Shannon Estuary are exposed or moderately exposed to wave action and subject to moderate tidal streams. Known sites are steeply sloping and show a good zonation down the shore. Well developed lichen zones and littoral reef communities offering a high species richness in the sublittoral fringe and strong populations of *Paracentrotus lividus* are found. The communities found are tolerant to sand scour and tidal streams. The infralittoral reefs range from sloping platforms with some vertical steps to ridged bedrock with gullies of sand between the ridges to ridged bedrock with boulders or a mixture of cobbles, gravel and sand. Kelp is very common to about 18m. Below this it becomes rare and the community is characterised by coralline crusts and red foliose algae.

Other coastal habitats that occur within the site include the following:

- stony beaches and bedrock shores - these shores support a typical zonation of seaweeds (*Fucus* spp., *Ascophyllum nodosum* and kelps).
- shingle beaches - the more stable areas of shingle support characteristic species such as Sea Beet, Sea Mayweed (*Matricaria maritima*), Sea Campion and Curled Dock (*Rumex crispus*).
- Sandbanks which are slightly covered by sea water at all times – there is a known occurrence of sand/gravel beds in the area from Kerry Head to Beal Head.
- sand dunes - a small area of sand dunes occurs at Beal Point. The dominant species is Marram Grass (*Ammophila arenaria*).

Flowing into the estuaries are a number of tidal rivers.

Freshwater rivers have been included in the site, most notably the Feale and Mulkear catchments, the Shannon from Killaloe to Limerick (along with some of its tributaries, including a short stretch of the Kilmastulla River), the Fergus up as far as Ennis, and the Cloon River. These systems are very different in character: the Shannon being broad, generally slow-flowing and naturally eutrophic; the Fergus being smaller and alkaline; while the narrow, fast-flowing Cloon is acid in nature. The Feale and Mulkear catchments exhibit all the aspects of a river from source to mouth. Semi-natural habitats, such as wet grassland, wet woodland and marsh occur by the rivers, however, improved grassland is most common. One grassland type of

particular conservation significance, *Molinia* meadows, occurs in several parts of the site and the examples at Worldsend on the River Shannon are especially noteworthy. Here are found areas of wet meadow dominated by rushes and sedges and supporting a diverse and species-rich vegetation, including such uncommon species as Blue-eyed Grass (*Sisyrinchium bermudiana*) and Pale Sedge (*Carex pallescens*).

Floating river vegetation characterised by species of Water-crowfoot (*Ranunculus* spp.), Pondweeds (*Potamogeton* spp.) and the moss *Fontinalis antipyretica* are present throughout the major river systems within the site. The rivers contain an interesting bryoflora with *Schistidium alpicola* var. *alpicola* recorded from in-stream boulders on the Bilboa, new to county Limerick.

Alluvial woodland occurs on the banks of the Shannon and on islands in the vicinity of the University of Limerick. The woodland is up to 50m wide on the banks and somewhat wider on the largest island. The most prominent woodland type is gallery woodland where White Willow (*Salix alba*) dominates the tree layer with occasional Alder (*Alnus glutinosa*). The shrub layer consists of various willow species with sally (*Salix cinerea* ssp. *oleifolia*) and what appear to be hybrids of *S. alba* x *S. viminalis*. The herbaceous layer consists of tall perennial herbs. A fringe of Bulrush (*Typha* sp.) occurs on the riverside of the woodland. On slightly higher ground above the wet woodland and on the raised embankment remnants of mixed oak-ash-alder woodland occur. These are poorly developed and contain numerous exotic species but locally there are signs that it is invading open grassland. Alder is the principal tree species with occasional Oak (*Quercus robur*), Elm (*Ulmus glabra*, *U. procera*), Hazel (*Corylus avellana*), Hawthorn (*Crataegus monogyna*) and the shrubs Guelder-rose (*Viburnum opulus*) and willows. The ground flora is species-rich.

Woodland is infrequent within the site, however Cahiracon Wood contains a strip of old Oak woodland. Sessile Oak (*Quercus petraea*) forms the canopy, with an understorey of Hazel and Holly (*Ilex aquifolium*). Great Wood-rush (*Luzula sylvatica*) dominates the ground flora. Less common species present include Great Horsetail (*Equisetum telmateia*) and Pendulous Sedge (*Carex pendula*).

In the low hills to the south of the Slievefelim mountains, the Cahernahallia River cuts a valley through the Upper Silurian rocks. For approximately 2km south of Cappagh Bridge at Knockanavar, the valley sides are wooded. The woodland consists of Birch (*Betula* spp.), Hazel, Oak, Rowan (*Sorbus aucuparia*), some Ash (*Fraxinus excelsior*) and Willow (*Salix* spp.). Most of the valley is not grazed by stock, and as a result the trees are regenerating well. The ground flora feature prominent Greater wood-rush and Bilberry (*Vaccinium myrtillus*) with a typical range of woodland herbs. Where there is more light available, Bracken (*Pteridium aquilinum*) features.

The valley sides of the Bilboa and Gortnageragh Rivers, on higher ground north east of Cappamore, support patches of semi-natural broadleaf woodland dominated by Ash, Hazel, Oak and Birch. There is a good scrub layer with Hawthorn, Willow, Holly and Blackthorn (*Prunus spinosa*) common. The herb layer in these woodlands is often open with a typically rich mixture of woodland herbs and ferns. Moss species diversity is high. The woodlands are ungrazed. The hazel is actively coppiced in places.

There is a small area of actively regenerating cut away raised bog at Ballyrorheen. It is situated approx. 5km north west of Cappamore Co. Limerick. The bog contains some wet areas with good moss (*Sphagnum*) cover. Species of particular interest include the Cranberry (*Vaccinium oxycoccos*) and the White Sedge (*Carex curta*) along with two other regionally rare mosses including *S. fimbriatum*. The site is being invaded by Birch (*Betula pubescens*) scrub woodland. Both commercial forestry and the spread of rhododendron has greatly reduced the overall value of the site.

A number of plant species that are Irish Red Data Book species occur within the site - several are protected under the Flora (Protection) Order, 1999:

- Triangular Club-rush (*Scirpus triquetrus*) - in Ireland this protected species is only found in the Shannon Estuary, where it borders creeks in the inner estuary.
- Opposite-leaved Pondweed (*Groenlandia densa*) - this protected pondweed is found in the Shannon where it passes through Limerick City.
- Meadow Barley (*Hordeum secalinum*) - this protected species is abundant in saltmarshes at Ringmoylan and Mantlehill.
- Hairy Violet (*Viola hirta*) - this protected violet occurs in the Askeaton/Foynes area.
- Golden Dock (*Rumex maritimus*) - noted as occurring in the River Fergus Estuary.
- Bearded Stonewort (*Chara canescens*) - a brackish water specialist found in Shannon Airport lagoon.
- Convergent Stonewort (*Chara conniven*) - presence in Shannon Airport Lagoon to be confirmed.

Overall, the Shannon and Fergus Estuaries support the largest numbers of wintering waterfowl in Ireland. The highest count in 1995-96 was 51,423 while in 1994-95 it was 62,701. Species listed on Annex I of the E.U. Birds Directive which contributed to these totals include: Great Northern Diver (3; 1994/95), Whooper Swan (201; 1995/96), Pale-bellied Brent Goose (246; 1995/96), Golden Plover (11,067; 1994/95) and Bar-tailed Godwit (476; 1995/96). In the past, three separate flocks of Greenland White-fronted Goose were regularly found but none were seen in 1993/94.

Other wintering waders and wildfowl present include Greylag Goose (216; 1995/96), Shelduck (1,060; 1995/96), Wigeon (5,976; 1995/96); Teal (2,319; 1995-96); Mallard (528; 1995/96), Pintail (45; 1995/96), Shoveler (84; 1995/96), Tufted Duck (272; 1995/96), Scaup (121; 1995/96), Ringed Plover (240; 1995/96), Grey Plover (750; 1995/96), Lapwing (24,581; 1995/96), Knot (800; 1995/96), Dunlin (20,100; 1995/96), Snipe (719; 1995/96), Black-tailed Godwit (1062; 1995/96), Curlew (1504; 1995/96), Redshank (3228; 1995/96), Greenshank (36; 1995/96) and Turnstone (107; 1995/96). A number of wintering gulls are also present, including Black-headed Gull (2,216; 1995/96), Common Gull (366; 1995/96) and Lesser Black-backed Gull (100; 1994/95). This is the most important coastal site in Ireland for a number of the waders including Lapwing, Dunlin, Snipe and Redshank. It also provides an important staging ground for species such as Black-tailed Godwit and Greenshank.



A number of species listed on Annex I of the E.U. Birds Directive breed within the site. These include Peregrine Falcon (2-3 pairs), Sandwich Tern (34 pairs on Rat Island, 1995), Common Tern (15 pairs: 2 on Sturamus Island and 13 on Rat Island, 1995), Chough (14-41 pairs, 1992) and Kingfisher. Other breeding birds of note include Kittiwake (690 pairs at Loop Head, 1987) and Guillemot (4010 individuals at Loop Head, 1987)

There is a resident population of Bottle-nosed Dolphin in the Shannon Estuary consisting of at least 56-68 animals (1996). This is the only known resident population of this E.U. Habitats Directive Annex II species in Ireland. Otter, a species also listed on Annex II of this directive, is commonly found on the site.

Five species of fish listed on Annex II of the E.U. Habitats Directive are found within the site. These are Sea Lamprey (*Petromyzon marinus*), Brook Lamprey (*Lampetra planeri*), River Lamprey (*Lampetra fluviatilis*), Twaite Shad (*Allosa fallax fallax*) and Salmon (*Salmo salar*). The three lampreys and Salmon have all been observed spawning in the lower Shannon or its tributaries. The Fergus is important in its lower reaches for spring salmon while the Mulkear catchment excels as a grilse fishery though spring fish are caught on the actual Mulkear River. The Feale is important for both types. Twaite Shad is not thought to spawn within the site. There are few other river systems in Ireland which contain all three species of Lamprey.

Two additional fish of note, listed in the Irish Red Data Book, also occur, namely Smelt (*Osmerus eperlanus*) and Pollan (*Coregonus autumnalis pollan*). Only the former has been observed spawning in the Shannon.

Freshwater Pearl-mussel (*Margaritifera margaritifera*), a species listed on Annex II of the E.U. Habitats Directive, occurs abundantly in parts of the Cloon River.

There is a wide range of landuses within the site. The most common use of the terrestrial parts is grazing by cattle and some areas have been damaged through over-grazing and poaching. Much of the land adjacent to the rivers and estuaries has been improved or reclaimed and is protected by embankments (especially along the Fergus Estuary). Further, reclamation continues to pose a threat as do flood relief works (e.g. dredging of rivers). Gravel extraction poses a major threat on the Feale.

In the past, Cord-grass (*Spartina* sp.) was planted to assist in land reclamation. This has spread widely, and may oust less vigorous colonisers of mud and may also reduce the area of mudflat available to feeding birds.

Domestic and industrial wastes are discharged into the Shannon, but water quality is generally satisfactory - except in the upper estuary, reflecting the sewage load from Limerick City. Analyses for trace metals suggest a relatively clean estuary with no influences by industrial discharges apparent. Further industrial development along the Shannon and water polluting operations are potential threats.

Fishing is a main tourist attraction on the Shannon and there are a large number of Angler Associations, some with a number of beats. Fishing stands and styles have been erected in places. The River Feale is a designated Salmonid Water under the

E.U. Freshwater Fish Directive. Other uses of the site include commercial angling, oyster farming, boating (including dolphin-watching trips) and shooting. Some of these may pose threats to the birds and dolphins through disturbance. Specific threats to the dolphins include underwater acoustic disturbance, entanglement in fishing gear and collisions with fast moving craft.

This site is of great ecological interest as it contains a high number of habitats and species listed on Annexes I and II of the E.U. Habitats Directive, including the priority habitat lagoon, the only known resident population of Bottle-nosed Dolphin in Ireland and all three Irish lamprey species. A good number of Red Data Book species are also present, perhaps most notably the thriving populations of Triangular Club-rush. A number of species listed on Annex I of the E.U. Birds Directive are also present, either wintering or breeding. Indeed, the Shannon and Fergus Estuaries form the largest estuarine complex in Ireland and support more wintering wildfowl and waders than any other site in the country. Most of the estuarine part of the site has been designated a Special Protection Area (SPA), under the E.U. Birds Directive, primarily to protect the large numbers of migratory birds present in winter.

6.10.2006

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

## SITE SYNOPSIS

**SITE NAME: RIVER SHANNON AND RIVER FERGUS ESTUARIES SPA**

**SITE CODE: 004077**

The estuaries of the River Shannon and River Fergus form the largest estuarine complex in Ireland. The site comprises all of the estuarine habitat west from Limerick City and south from Ennis, extending west as far as Killadysert and Foynes on the north and south shores respectively of the River Shannon (a distance of some 25 km from east to west). Also included are several areas in the outer Shannon estuary, notably Clonderalaw Bay and Poulnasherry Bay, as well as the intertidal areas on the south shore of the Shannon between Tarbert and Beal Point.

The site has vast expanses of intertidal flats. The main macro-invertebrate community present is a *Macoma-Scrobicularia-Nereis* community which provides a rich food resource for the wintering birds. Other species occurring include Common Cockle (*Cerastoderma edule*), Lugworm (*Arenicola marina*), the polychaete *Nephtys hombergii*, the gastropod *Hydrobia ulvae* and the crustacean *Corophium volutator*. Eelgrass (*Zostera* spp.) is present in places, along with green algae (e.g. *Ulva* spp. and *Enteromorpha* spp.). Salt marsh vegetation frequently fringes the mudflats and this provides important high tide roost areas for the wintering birds. Characteristic species occurring include Common Saltmarsh-grass (*Puccinellia maritima*), Sea Aster (*Aster tripolium*), Thrift (*Armeria maritima*), Sea-milkwort (*Glaux maritima*), Sea Plantain (*Plantago maritima*), Red Fescue (*Festuca rubra*) and Saltmarsh Rush (*Juncus gerardi*). In the innermost parts of the estuaries, the tidal channels or creeks are fringed with species such as Common Reed (*Phragmites australis*) and club-rushes (*Scirpus maritimus*, *S. lacustris* subsp. *tabernaemontani*). Also found is the nationally rare Triangular Club-rush (*Scirpus triqueter*). Elsewhere in the site the shoreline comprises stony or shingle beaches.

The site is the most important coastal wetland site in the country and regularly supports in excess of 50,000 wintering waterfowl (mean of 59,183 for the 4 seasons 1996-97 to 1999/00), a concentration easily of international importance. The site has internationally important populations of Dunlin (14,987), Black-tailed Godwit (706) and Redshank (1,983) - all figures are average peaks for 3 of the 5 seasons in the 1995/96-1999/00 period. A further 16 species have populations of national importance, i.e. Cormorant (148), Whooper Swan (141), Greylag Goose (88), Shelduck (895), Wigeon (3,025), Teal (1,558), Pintail (40), Shoveler (56), Scaup (76), Golden Plover (4,073), Grey Plover (564), Lapwing (13,007), Knot (686), Bar-tailed Godwit (481), Curlew (1,231) and Greenshank (33). The site is among the most important in the country for several of these species, notably Dunlin (11% of national total), Grey Plover (7.5% of total), Lapwing (6.5% of total), Redshank (6% of total) and Shelduck (6.0% of total). The site is also used by Oystercatcher (363), Ringed Plover (70), Brent Goose (135), Great Crested Grebe (47), Red-breasted Merganser (14), Mallard (247), Turnstone (71), Mute Swan (54), Grey Heron (25), Black-headed Gull (1,233) and Common Gull (194).

The Shannon / Fergus system was formerly frequented by a Greenland White-fronted Goose population but this declined during the 1980s and 1990s and the birds now appear to have abandoned the area. The site provides both feeding and roosting areas for the wintering birds. Habitat quality for most of the estuarine habitats is good. Some species, particularly Whooper Swan and Greylag Goose, utilise areas outside of the site for feeding.

Apart from the wintering birds, large numbers of some species also pass through the site whilst on migration in spring and/or autumn. Regular species include Black-tailed Godwit, Whimbrel and Greenshank.

Much of the land adjacent to the rivers and estuaries has been reclaimed and improved for agriculture and is protected by embankments (especially along the River Fergus estuary). Further reclamation, especially near to the urbanised and industrial areas continues to pose a threat. The site receives pollution from several sources, including industry and agriculture, but it is not known if this has any significant impacts on the wintering birds. Aquaculture occurs in some areas of the site – future increases in this activity could cause disturbance to the habitats and the associated birds. Common Cord-grass (*Spartina anglica*) is well-established and may threaten some of the estuarine habitats. Some disturbance occurs from boating activities.

This site is of great ornithological interest, being of international importance on account of the numbers of wintering birds it supports. It also supports internationally important numbers of three species, i.e. Dunlin, Black-tailed Godwit and Redshank. In addition, there are 16 species that have populations of national importance. For several of the bird species, it is the top site in the country. Also of note is that three of the species which occur regularly are listed on Annex I of the E.U. Birds Directive, i.e. Whooper Swan, Golden Plover and Bar-tailed Godwit. The site is most effectively censused from the air and this is carried out in most winters.

# APPENDIX 3

For inspection purposes only.  
Consent of copyright owner required for any other use.

# National Parks and Wildlife Service

## Conservation Objectives Series

### Lower River Shannon SAC 002165

For inspection purposes only.  
Consent of copyright owner required for any other use.



**An Roinn**  
**Ealaíon, Oidhreachta agus Gaeltachta**  
**Department of**  
**Arts, Heritage and the Gaeltacht**



**National Parks and Wildlife Service,  
Department of Arts, Heritage and the Gaeltacht,  
7 Ely Place, Dublin 2, Ireland.  
Web: [www.npws.ie](http://www.npws.ie)  
E-mail: [nature.conservation@ahg.gov.ie](mailto:nature.conservation@ahg.gov.ie)**

**Citation:**

NPWS (2012) Conservation Objectives: Lower River Shannon SAC 002165. Version 1.0. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht.

**Series Editors: Rebecca Jeffrey & Naomi Kingston  
ISSN 2009-4086**

## Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

### Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.



## Qualifying Interests

\* indicates a priority habitat under the Habitats Directive

### 002165 Lower River Shannon SAC

- 1029 Freshwater Pearl Mussel *Margaritifera margaritifera*
- 1095 Sea Lamprey *Petromyzon marinus*
- 1096 Brook Lamprey *Lampetra planeri*
- 1099 River Lamprey *Lampetra fluviatilis*
- 1106 Atlantic Salmon *Salmo salar* (only in fresh water)
- 1110 Sandbanks which are slightly covered by sea water all the time
- 1130 Estuaries
- 1140 Mudflats and sandflats not covered by seawater at low tide
- 1150 \*Coastal lagoons
- 1160 Large shallow inlets and bays
- 1170 Reefs
- 1220 Perennial vegetation of stony banks
- 1230 Vegetated sea cliffs of the Atlantic and Baltic coasts
- 1310 *Salicornia* and other annuals colonizing mud and sand
- 1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritima*)
- 1349 Bottlenose Dolphin *Tursiops truncatus*
- 1355 Otter *Lutra lutra*
- 1410 Mediterranean salt meadows (*Juncetalia maritimi*)
- 3260 Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitriche-Batrachion* vegetation
- 6410 *Molinia* meadows on calcareous, peaty or clayey-silt-laden soils (*Molinion caeruleae*)
- 91E0 \*Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (*Alno-Padion*, *Alnion incanae*, *Salicion albae*)

Please note that this SAC overlaps with River Shannon and River Fergus Estuaries SPA (004077), Loop Head SPA (004119), Stack's to Mullaghareirk Mountains, West Limerick Hills and Mount Eagle SPA (004161), Slievefelim to Silvermines Mountains SPA (004165) and Kerry Head SPA (004189). It is also adjacent to Clare Glen SAC (00930). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping and adjacent sites as appropriate.

## Supporting documents, relevant reports & publications (listed by date)

Supporting documents, NPWS reports and publications are available for download from: [www.npws.ie/Publications](http://www.npws.ie/Publications)

---

**Title:** Aspects of brook lamprey (*Lampetra planeri* Bloch) spawning in Irish waters

**Year:** in press

**Author:** Rooney, S.M.; O’Gorman, N.M.; Green, F.; King, J.J.

**Series:** Biology and Environment

---

**Title:** Lower River Shannon SAC (002170): Conservation objectives supporting document - Coastal lagoons [Version 1]

**Year:** 2012

**Author:** NPWS

**Series:** Unpublished Report to NPWS

---

**Title:** Lower River Shannon SAC (002170): Conservation objectives supporting document - Marine habitats and species [Version 1]

**Year:** 2012

**Author:** NPWS

**Series:** Unpublished Report to NPWS

---

**Title:** Lower River Shannon SAC (002170): Conservation objectives supporting document - Coastal habitats [Version 1]

**Year:** 2012

**Author:** NPWS

**Series:** Unpublished Report to NPWS

---

**Title:** Lower River Shannon SAC (002170): Conservation objectives supporting document - Woodland habitats [Version 1]

**Year:** 2012

**Author:** NPWS

**Series:** Unpublished Report to NPWS

---

**Title:** Lower River Shannon SAC (002170): Conservation objectives supporting document - Water courses of plain to montane levels with the *Ranunculon fluitantis* and *Callitricho-Batrachion* vegetation [Version 1]

**Year:** 2012

**Author:** NPWS

**Series:** Unpublished Report to NPWS

---

**Title:** Intertidal Hard and Soft Bottom Investigations in Lower River Shannon cSAC (Site Code: IE002165)/Shannon Fergus Estuary SPA (Site Code: IE004077)

**Year:** 2011c

**Author:** Aquafact

**Series:** Unpublished Report to NPWS

---

**Title:** Reef Investigations in Lower River Shannon cSAC (cSAC Site Code: IE002165)

**Year:** 2011b

**Author:** Aquafact

**Series:** Unpublished Report to NPWS

---

---

**Title:** Subtidal Benthic Investigations in Lower River Shannon cSAC (cSAC Site Code: IE002165)  
**Year:** 2011a  
**Author:** Aquafact  
**Series:** Unpublished Report to NPWS

---

**Title:** National survey and assessment of the conservation status of Irish sea cliffs  
**Year:** 2011  
**Author:** Barron, S.J.; Delaney, A.; Perrin, P.M.; Martin, J.; O'Neill, F.  
**Series:** Irish Wildlife Manuals No. 53

---

**Title:** Comparison of field- and GIS-based assessments of barriers to Atlantic salmon migration: a case study in the Nore Catchment, Republic of Ireland  
**Year:** 2011  
**Author:** Gargan, P. G.; Roche, W. K.; Keane, S.; King, J.J.; Cullagh, A.; Mills, P.; O'Keeffe, J.  
**Series:** J. Appl. Ichthyol. 27 (Suppl. 3), 66–72

---

**Title:** Fine-scale population genetic structuring of bottlenose dolphins in Irish coastal waters  
**Year:** 2011  
**Author:** Mirimin, L.; Miller, R.; Dillane, E.; Berrow, S.D.; Ingram, S.; Cross, T.F.; Rogan, E.  
**Series:** Animal Conservation 2011: 1–12

---

**Title:** The use of Cork Harbour by bottlenose dolphins (*Tursiops truncatus* (Montagu, 1821))  
**Year:** 2011  
**Author:** Ryan, C.; Cross, T.F.; Rogan, E.  
**Series:** Irish Naturalists' Journal 31(1): 1-9

---

**Title:** Irish cetacean review (2000-2009)  
**Year:** 2010  
**Author:** Berrow, S.D.; Whooley, P.; O'Connell, M.; Wall, D.  
**Series:** Irish Whale and Dolphin Group

---

**Title:** Bottlenose Dolphin SAC Survey 2010  
**Year:** 2010  
**Author:** Berrow, S.D.; O'Brien, J.; Groth, L.; Foley, A.; Voigt, K.  
**Series:** Unpublished Report to NPWS

---

**Title:** Otter tracking study of Roaringwater Bay  
**Year:** 2010  
**Author:** De Jongh, A.; O'Neill, L.  
**Series:** Unpublished Draft Report to NPWS

---

**Title:** Second Draft Cloon (Shannon Estuary) Freshwater Pearl Mussel Sub-basin Management Plan (2009-2015)  
**Year:** 2010  
**Author:** DEHLG  
**Series:** Unpublished Report to NPWS

---

---

**Title:** Social structure within the bottlenose dolphin (*Tursiops truncatus*) population in the Shannon Estuary, Ireland

**Year:** 2010

**Author:** Foley, A.; McGrath, D.; Berrow, S.D.; Gerritsen, H.

**Series:** Aquatic Mammals 36(4): 372-381

---

**Title:** Irish Semi-natural Grasslands Survey. Annual report no. 3: Counties Donegal, Dublin, Kildare & Sligo

**Year:** 2010

**Author:** O'Neill, F.H.; Martin, J.R.; Devaney, F.M.; McNutt, K.E.; Perrin, P.M.; Delaney, A.

**Series:** Unpublished Report to NPWS

---

**Title:** A provisional inventory of ancient and long-established woodland in Ireland

**Year:** 2010

**Author:** Perrin, P.M.; Daly, O.H.

**Series:** Irish Wildlife Manuals No. 46

---

**Title:** Monitoring and Assessment of Irish Lagoons for the purpose of the EU Water Framework Directive

**Year:** 2010

**Author:** Roden, C.M.; Oliver, G.

**Series:** EPA

---

**Title:** Report of the standing scientific committee to the DCENR. The status of Irish salmon stocks in 2010 and precautionary catch advice for 2011

**Year:** 2010

**Author:** SSC

**Series:** Unpublished Report to DCENR

---

**Title:** The European Communities Environmental Objectives (Freshwater Pearl Mussel) Regulations 2009. [S.I. 296 of 2009]

**Year:** 2009b

**Author:** Government of Ireland

**Series:** Irish Statute Book

---

**Title:** The European Communities Environmental Objectives (Surface Water) Regulations 2009. [S.I. 272 of 2009]

**Year:** 2009a

**Author:** Government of Ireland

**Series:** Irish Statute Book

---

**Title:** Winter distribution of bottle-nosed dolphins (*Tursiops truncatus* (Montagu)) in the inner Shannon Estuary

**Year:** 2009

**Author:** Berrow, S.D.

**Series:** Irish Naturalists' Journal 30(1): 35-39

---

**Title:** Towards a bottlenose dolphin whistle ethogram from the Shannon Estuary, Ireland

**Year:** 2009

**Author:** Hickey, R.; Berrow, S.D.; Goold, J.

**Series:** Biology and Environment: Proceedings of the Royal Irish Academy 109B (2), 89-94

---

---

**Title:** Saltmarsh Monitoring Report 2007-2008  
**Year:** 2009  
**Author:** McCorry, M.; Ryle, T.  
**Series:** Unpublished Report to NPWS

---

**Title:** Cetaceans in Irish waters: A review of recent research  
**Year:** 2009  
**Author:** O'Brien, J.; Berrow, S.D.; McGrath, D.; Evans, P.G.H.  
**Series:** Biology and Environment: Proceedings of the Royal Irish Academy 109B (2): 63-88

---

**Title:** A note on long-distance matches of bottlenose dolphins (*Tursiops truncatus*) around the Irish coast using photoidentification  
**Year:** 2009  
**Author:** O'Brien, J.; Berrow, S.D.; Ryan, C.; McGrath, D.; O'Connor, I.; Pesante, G.; Burrows, G.; Massett, N.; Klotzer, V.; Whooley, P.  
**Series:** Journal Cetacean Res. Mgmt. 11: 69–74

---

**Title:** An updated population status report for bottlenose dolphins using the Lower River Shannon SAC in 2008  
**Year:** 2008  
**Author:** Englund, A.; Ingram, S.; Rogan, E.  
**Series:** Unpublished Report to NPWS

---

**Title:** National Survey of Native Woodlands 2003-2008  
**Year:** 2008  
**Author:** Perrin, P.; Martin, J.; Barron, S.; O'Neill, E.; McNutt, K.; Delaney, A.  
**Series:** Unpublished Report to NPWS

---

**Title:** Rapid Assessment of *Margaritifera margaritifera* (L.) populations in Ireland: Rivers assessed in 2007  
**Year:** 2008  
**Author:** Ross, E.D.  
**Series:** Unpublished Report to NPWS

---

**Title:** Marine surveys of two Irish sandbank cSACs  
**Year:** 2007  
**Author:** Aquafact  
**Series:** Unpublished Report to NPWS

---

**Title:** Population status report for bottlenose dolphins using the Lower River Shannon SAC, 2006-2007  
**Year:** 2007  
**Author:** Englund, A.; Ingram, S.; Rogan, E.  
**Series:** Unpublished Report to NPWS

---

**Title:** Evolutionary history of lamprey paired species *Lampetra fluviatilis* (L.) and *Lampetra planeri* (Bloch) as inferred from mitochondrial DNA variation  
**Year:** 2007  
**Author:** Espanhol, R.; Almeida, P.R.; Alves, M.J.  
**Series:** Molecular Ecology 16, 1909-1924

---

---

**Title:** Supporting documentation for the Habitats Directive Conservation Status Assessment - backing documents, Article 17 forms and supporting maps

**Year:** 2007

**Author:** NPWS

**Series:** Unpublished Report to NPWS

---

**Title:** A Survey of Juvenile Lamprey Populations in the Corrib and Suir Catchments

**Year:** 2007

**Author:** O'Connor, W.

**Series:** Irish Wildlife Manuals No. 26

---

**Title:** Inventory of Irish coastal lagoons

**Year:** 2007

**Author:** Oliver, G.

**Series:** Unpublished Report to NPWS

---

**Title:** Using T-PODs to investigate the echolocation of coastal bottlenose dolphins

**Year:** 2007

**Author:** Philpott, E.; Englund, A.; Ingram, S.; Rogan, E.

**Series:** Journal of Marine Biological Association, UK. 87: 11-17

---

**Title:** Otter Survey of Ireland 2004/2005

**Year:** 2006

**Author:** Bailey, M.; Rochford, J.

**Series:** Irish Wildlife Manuals No. 23

---

**Title:** Whistle Production by Bottlenose Dolphins *Tursiops truncatus* in the Shannon Estuary

**Year:** 2006

**Author:** Berrow, S.D.; O'Brien, J.; Holmes, B.

**Series:** Irish Naturalists' Journal. 28(5): 208-213

---

**Title:** The status of host fish populations and fish species richness in European freshwater pearl mussel (*Margaritifera margaritifera*) streams

**Year:** 2006

**Author:** Geist, J.; Porkka, M.; Kuehn, R.

**Series:** Aquatic Conservation: Marine and Freshwater Ecosystems 16, 251–266

---

**Title:** Otters - ecology, behaviour and conservation

**Year:** 2006

**Author:** Kruuk, H.

**Series:** Oxford University Press

---

**Title:** A survey of rare and scarce vascular plants in County Limerick

**Year:** 2006

**Author:** Reynolds, S.; Conaghan, J.; Fuller, J.

**Series:** Unpublished Report to NPWS

---

- Title:** National Inventory of sea cliffs and coastal heaths  
**Year:** 2005  
**Author:** Browne, A.  
**Series:** Unpublished Report to NPWS
- 
- Title:** Developing sustainable whalewatching in the Shannon estuary  
**Year:** 2003  
**Author:** Berrow, S.D.  
**Series:** p198-203; In Marine Ecotourism: Issues and Experiences. Garrod, B and Wilson. J. (Eds.) Channel View Publications
- 
- Title:** Identifying lamprey. A field key for sea, river and brook lamprey  
**Year:** 2003  
**Author:** Gardiner, R.  
**Series:** Conserving Natura 2000 rivers, Conservation techniques No. 4. English Nature, Peterborough
- 
- Title:** Monitoring the river, sea and brook lamprey, *Lampetra fluviatilis*, *L. planeri* and *Petromyzon marinus*  
**Year:** 2003  
**Author:** Harvey, J.; Cowx, I.  
**Series:** Conserving Natura 2000 Rivers Monitoring Series No. 5. English Nature, Peterborough
- 
- Title:** Bottlenose dolphins (*Tursiops truncatus*) in the Shannon Estuary and selected areas of the west-coast of Ireland  
**Year:** 2003  
**Author:** Ingram, S.; Rogan, E.  
**Series:** Unpublished Report to NPWS
- 
- Title:** The ecology of seabirds and marine mammals in a fluctuating marine environment  
**Year:** 2003  
**Author:** Rogan, E.; Kelly, T.; Ingram, S.; Roycroft, D.  
**Series:** Unpublished Report to Higher Education Authority of Ireland
- 
- Title:** Irish Whale and Dolphin Group cetacean sighting review (1991-2001)  
**Year:** 2002  
**Author:** Berrow, S.D.; Whooley, P.; Ferriss, S.  
**Series:** Irish Whale and Dolphin Group
- 
- Title:** Organochlorine concentrations in resident bottlenose dolphins (*Tursiops truncatus*) in the Shannon estuary, Ireland  
**Year:** 2002  
**Author:** Berrow, S.D.; McHugh, B.; Glynn, D.; McGovern, E.; Parsons, K.; Baird, R.W.; Hooker, S.D.  
**Series:** Marine Pollution Bulletin 44: 1296-1313
- 
- Title:** Identifying critical areas and habitat preferences of bottlenose dolphins (*Tursiops truncatus*)  
**Year:** 2002  
**Author:** Ingram, S.; Rogan, E.  
**Series:** Marine Ecology Progress Series 244: 247-255
-

- 
- Title:** Reversing the habitat fragmentation of British woodlands  
**Year:** 2002  
**Author:** Peterken, G.  
**Series:** WWF-UK, London
- 
- Title:** An extensive survey of bottlenose dolphins (*Tursiops truncatus*) on the west coast of Ireland  
**Year:** 2001  
**Author:** Ingram, S.; Englund, A.; Rogan, E.  
**Series:** Unpublished Report to the Heritage Council
- 
- Title:** The ecology and conservation of bottlenose dolphins in the Shannon Estuary, Ireland  
**Year:** 2000  
**Author:** Ingram, S.  
**Series:** Unpublished PhD thesis, University College Cork
- 
- Title:** A survey of bottlenose dolphins (*Tursiops truncatus*) in the Shannon Estuary  
**Year:** 2000  
**Author:** Rogan, E.; Ingram, S.; Holmes, B.; O'Flanagan, C.  
**Series:** Marine Institute Marine Resource Series No. 9
- 
- Title:** Tour boats and dolphins: A note on quantifying the activities of whale watching boats in the Shannon estuary, Ireland  
**Year:** 1999  
**Author:** Berrow, S.D.; Holmes, B.  
**Series:** Journal of Cetacean Research and Management 1(2): 199-200
- 
- Title:** Diet of Otters *Lutra lutra* on Inishmore, Aran Islands, west coast of Ireland  
**Year:** 1999  
**Author:** Kingston, S.; O'Connell, M.; Fairley, J.S.  
**Series:** Biol & Environ Proc R Ir Acad B 99B:173-182
- 
- Title:** National Shingle Beach Survey of Ireland 1999  
**Year:** 1999  
**Author:** Moore, D.; Wilson, F.  
**Series:** Unpublished Report to NPWS
- 
- Title:** The saltmarshes of Ireland: an inventory and account of their geographical variation  
**Year:** 1998  
**Author:** Curtis, T.G.F.; Sheehy-Skeffington, M.J.  
**Series:** Biology and Environment, Proceedings of the Royal Irish Academy 98B: 87-104
- 
- Title:** A survey of intertidal sediment biotopes in estuaries in Ireland  
**Year:** 1997  
**Author:** Falvey, J.P.; Costello, M.J.; Dempsey, S.  
**Series:** Unpublished Report
-



---

**Title:** Distribution and Abundance of Bottle-nosed Dolphins *Tursiops truncatus* (Montagu) in the Shannon Estuary, Ireland

**Year:** 1996

**Author:** Berrow, S.D.; Holmes, B.; Kiely, O.

**Series:** Biology and Environment: Proceedings of the Royal Irish Academy 96B (1), 1-9

---

**Title:** The spatial organization of otters (*Lutra lutra*) in Shetland

**Year:** 1991

**Author:** Kruuk, H.; Moorhouse, A.

**Series:** J. Zool, 224: 41-57

---

**Title:** Otter survey of Ireland

**Year:** 1982

**Author:** Chapman, P.J.; Chapman, L.L.

**Series:** Unpublished Report to Vincent Wildlife Trust

---

For inspection purposes only.  
Consent of copyright owner required for any other use.

## Spatial data sources

<b>Year:</b>	Interpolated 2012
<b>Title:</b>	Sandbank Survey 2007
<b>GIS operations:</b>	Clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	1110 (map 3)
<b>Year:</b>	Interpolated 2012
<b>Title:</b>	Sandbank survey 2007; subtidal benthic survey 2010; reef survey 2010; intertidal hard and soft bottom survey 2010
<b>GIS operations:</b>	Polygon feature classes from marine community types base data sub-divided based on interpolation of marine survey data. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	Marine community types, 1110, 1140, 1170 (maps 3, 5, 8, 9)
<b>Year:</b>	2010
<b>Title:</b>	EPA WFD transitional waterbody data
<b>GIS operations:</b>	Clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	1130 (map 4)
<b>Year:</b>	Revision 2011
<b>Title:</b>	Inventory of Irish Coastal Lagoons. Version 3
<b>GIS operations:</b>	Clipped to SAC boundary
<b>Used for:</b>	1150 (map 6)
<b>Year:</b>	2005
<b>Title:</b>	OSi Discovery series vector data
<b>GIS operations:</b>	High Water Mark (HWM) polyline feature class converted into polygon feature class; clipped to SAC boundary. EPA WFD transitional waterbody data erased from extent. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	1160 (map 7)
<b>Year:</b>	2005
<b>Title:</b>	OSi Discovery series vector data
<b>GIS operations:</b>	High water mark (HWM) and low water mark (LWM) polyline feature classes converted into polygon feature classes and combined; EU Annex I Saltmarsh and Coastal data erased out if present
<b>Used for:</b>	Marine community types base data (map 9)
<b>Year:</b>	Revision 2012
<b>Title:</b>	National Shingle Beach Survey
<b>GIS operations:</b>	Clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	1220 (map 10)
<b>Year:</b>	2011
<b>Title:</b>	National Survey and assessment of the conservation status of Irish sea cliffs
<b>GIS operations:</b>	Clipped to SAC boundary
<b>Used for:</b>	1230 (map 11)

<b>Year:</b>	Revision 2010
<b>Title:</b>	Saltmarsh Monitoring Project 2007-2008. Version 1
<b>GIS operations:</b>	QIs selected; clipped to SAC boundary; overlapping regions with Coastal CO data investigated and resolved with expert opinion used
<b>Used for:</b>	1310, 1330, 1410 (map 12)
<b>Year:</b>	Derived 2012
<b>Title:</b>	Internal NPWS files
<b>GIS operations:</b>	Dataset created from spatial references supplied by NPWS experts. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	3260 (map 13)
<b>Year:</b>	Revision 2010
<b>Title:</b>	National Survey of Native Woodlands 2003-2008. Version 1
<b>GIS operations:</b>	QIs selected; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	91E0 (map 14)
<b>Year:</b>	2012
<b>Title:</b>	NPWS rare and threatened species database
<b>GIS operations:</b>	Dataset created from spatial references in database records. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	1029 (map 15)
<b>Year:</b>	Revision 2012
<b>Title:</b>	Margaritifera Sensitive Areas data
<b>GIS operations:</b>	Relevant catchment boundaries identified. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	1029 (map 15)
<b>Year:</b>	2005
<b>Title:</b>	OSi Discovery series vector data
<b>GIS operations:</b>	Low Water Mark (LWM) polyline feature class converted into polygon feature class; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	1349 (map 16)
<b>Year:</b>	2005
<b>Title:</b>	OSi Discovery series vector data
<b>GIS operations:</b>	Creation of an 80m buffer on the marine side of the high water mark (HWM); creation of a 10m buffer on the terrestrial side of the HWM; combination of 80m and 10m HWM buffer datasets; creation of a 10m buffer on the terrestrial side of the river banks data; creation of 20m buffer applied to canal centreline data. These datasets are combined with the derived EPA WFD Waterbodies data and Coastal Lagoon data for the 1355 CO. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising. Creation of 250m buffer on marine side of HWM to highlight potential commuting points
<b>Used for:</b>	1355 (map 17)

---

<b>Year:</b>	2010
<b>Title:</b>	EPA WFD Waterbodies data
<b>GIS operations:</b>	Creation of a 20m buffer applied to river and stream centreline data; creation of 80m buffer on the aquatic side of lake data; creation of 10m buffer on the terrestrial side of lake data. These datasets are combined with the derived OSi data and Coastal Lagoon data for the 1355 CO. Overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	1355 (no map)

---

<b>Year:</b>	Revision 2011
<b>Title:</b>	Inventory of Irish Coastal Lagoons. Version 3
<b>GIS operations:</b>	Creation of 80m buffer on the aquatic side of lagoon data; creation of 10m buffer on the terrestrial side of lagoon data. These datasets are combined with the derived OSi data and EPA WFD Waterbodies data for the 1355 CO. Overlapping regions are investigated and resolved; resulting dataset clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising
<b>Used for:</b>	1355 (no map)

---

For inspection purposes only.  
Consent of copyright owner required for any other use.

**Conservation objectives for: Lower River Shannon SAC [002165]**

**1029 Freshwater Pearl Mussel *Margaritifera margaritifera***

To restore the favourable conservation condition of Freshwater Pearl Mussel in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Kilometres	Maintain at 7km. See map 15	This conservation objective applies to the freshwater pearl mussel population in the Cloon River, Co. Clare only (see also the European Communities Environmental Objectives (Freshwater Pearl Mussel) Regulations 2009 (Government of Ireland, 2009b)). The Cloon population is confined to the main channel and is distributed from Croany Bridge to approx. 1.5km upstream of Clonderalaw Bridge (Ross, 2008; DEHLG, 2010)
Population size	Number of adult mussels	Restore to 10,000 adult mussels	The Cloon population was estimated as less than 10,000 in 2009 (DEHLG, 2010)
Population structure: recruitment	Percentage per size class	Restore to least 20% of population no more than 65mm in length; and at least 5% of population no more than 30mm in length	Mussels of no more than 65mm are considered 'young mussels' and may be found buried in the substratum and/or beneath adult mussels. Mussels of no more than 30mm are 'juvenile mussels' and are always buried in the substratum. No juvenile or young mussels were found in the Cloon in 2007, with the smallest mussel measuring 80.3mm (Ross, 2008). A single 'young mussel' measuring 61.3mm was recorded in 2009 (DEHLG, 2010)
Population structure: adult mortality	Percentage	No more than 5% decline from previous number of live adults counted; dead shells less than 1% of the adult population and scattered in distribution	5% is considered the cut-off between the combined errors associated with natural fluctuations and sampling methods and evidence of true population decline. 1% of dead shells is considered to be indicative of natural losses. The Cloon failed the target for dead shells in 2009, with 31% dead shells across the single transect counted. There were no previous data on the number of live adults (DEHLG, 2010)

For inspection purposes only. Consent of copyright owner required for any other use.

1029 Freshwater Pearl Mussel *Margaritifera margaritifera*

To restore the favourable conservation condition of Freshwater Pearl Mussel in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat extent	Kilometres	Restore suitable habitat in more than 3.3km (see map 15) and any additional stretches necessary for salmonid spawning	The species' habitat covers stretches of a short coastal river; and is a combination of 1) the area of habitat adult and juvenile mussels can occupy and 2) the area of spawning and nursery habitats the host fish can occupy. Fish nursery habitat typically overlaps with mussel habitat. Fish spawning habitat is generally adjacent to mussel habitat, but may lie upstream of the generalised mussel distribution. Only those salmonid spawning areas that could regularly contribute juvenile fish to the areas occupied by adult mussels should be considered. The availability of mussel habitat and fish spawning and nursery habitats are determined by flow and substratum conditions. The habitat for the species is currently unsuitable for the survival of adult mussels or the recruitment of juveniles (DEHLG, 2010). The target is based on the stretches of river identified, from a combination of dedicated survey and incidental records, as having habitat for the species
Water quality: macroinvertebrate and phytobenthos (diatoms)	Ecological quality ratio (EQR)	Restore water quality-macroinvertebrates: EQR greater than 0.90; Phytobenthos: EQR greater than 0.93	These EQRs correspond to high ecological status for these two Water Framework Directive biological quality elements. They represent high water quality with very low nutrient concentrations (oligotrophic conditions). The habitat in the Cloon failed both standards during 2009 sampling for the Sub-basin Management Plans (DEHLG, 2010). See also The European Communities Environmental Objectives (Surface Water) Regulations 2009 (Government of Ireland, 2009a)
Substratum quality: filamentous algae (macroalgae), macrophytes (rooted higher plants)	Percentage	Restore substratum quality-filamentous algae: absent or trace (<5%); macrophytes: absent or trace (<5%)	The habitat in the Cloon failed both standards during 2009 sampling for the Sub-basin Management Plans, with cover abundance values of up to 50% recorded for filamentous algae and 80% for macrophytes (DEHLG, 2010). Recruitment of juvenile mussels is being prevented by the poor quality of the river substrata

Consent of copyright owner required for any other use.

1029 Freshwater Pearl Mussel *Margaritifera margaritifera*

To restore the favourable conservation condition of Freshwater Pearl Mussel in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Substratum quality: sediment	Occurrence	Restore substratum quality-stable cobble and gravel substrate with very little fine material; no artificially elevated levels of fine sediment	The habitat for the species is currently unsuitable for the recruitment of juveniles owing to sedimentation of the substratum. In many locations, it is also unsuitable for the survival of adult mussels (DEHLG, 2010). Significant sedimentation has been recorded during all recent mussel monitoring surveys (Ross, 2008; DEHLG, 2010). Recruitment of juvenile mussels is being prevented by the poor quality of the river substrate
Substratum quality: oxygen availability	Redox potential	Restore to no more than 20% decline from water column to 5cm depth in substrate	Differences in redox potential between the water column and the substrate correlate with differences in oxygen levels. Juvenile mussels require full oxygenation while buried in gravel. In suitable habitat, there should be very little loss of redox potential between the water column and underlying gravels. Redox potential measurements in 2009 yielded losses of 32.3 - 43.5% (average of 39%) at 5cm depth (DEHLG, 2010)
Hydrological regime: flow variability	Metres per second	Restore appropriate hydrological regimes	The availability of suitable freshwater pearl mussel habitat is largely determined by flow (catchment geology being the other important factor). In order to restore the habitat for the species, flow variability over the annual cycle must be such that: 1) high flows can wash fine sediments from the substratum, 2) low flows do not exacerbate the deposition of fines and 3) low flows do not cause stress to mussels in terms of exposure, water temperatures, food availability or aspects of the reproductive cycle

For inspection purposes only. Consent of copyright owner required for any other use.

1029 Freshwater Pearl Mussel *Margaritifera margaritifera*

To restore the favourable conservation condition of Freshwater Pearl Mussel in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Host fish	Number	Maintain sufficient juvenile salmonids to host glochidial larvae	Salmonid fish are host to the larval form of the freshwater pearl mussel and, thus, they are essential to the completion of the life cycle. 0+ and 1+ fish are typically used, both because of the habitat overlaps and the development of immunity with age in the fish. Fish presence is considered sufficient, as higher densities and biomass of fish are indicative of enriched conditions in mussel rivers. Geist et al. (2006) found that higher densities of host fish coincided with eutrophication, poor substrate quality for pearl mussels and a lack of pearl mussel recruitment, while significantly lower densities and biomass of host fish were associated with high numbers of juvenile mussels. Fish movement patterns must be such that 0+ fish in the vicinity of the mussel habitat remain in the mussel habitat until their 1+ summer. No fish stocking should occur within the mussel habitat, nor any works that may change the salmonid balance or residency time. The Cloon freshwater pearl mussel population appears to favour native brown trout, with 17.2% of 1+ and older trout caught in 2009 hosting glochidia (DEHLG, 2010). Therefore, it is particularly important that trout are not out-competed by stocked fish

For inspection purposes only.  
Consent of copyright owner required for any other use.



**Conservation objectives for: Lower River Shannon SAC [002165]**

**1095 Sea Lamprey *Petromyzon marinus***

**To restore the favourable conservation condition of Sea Lamprey in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Distribution: extent of anadromy	% of river accessible	Greater than 75% of main stem length of rivers accessible from estuary	Artificial barriers can block or cause difficulties to lampreys' upstream migration, thereby limiting the species to lower stretches and restricting access to spawning areas. See Gargan et al. (2011). Specific barriers serve to constrain the up-river migration of sea lamprey. The upper extent of the SAC in the R. Fergus is delineated by a barrier to migration. Barriers are also present in the Mulkear and Feale
Population structure of juveniles	Number of age/size groups	At least three age/size groups present	Attribute and target based on data from Harvey and Cowx (2003) and O'Connor (2007)
Juvenile density in fine sediment	Juveniles/m <sup>2</sup>	Juvenile density at least 1/m <sup>2</sup>	Juveniles burrow in areas of fine sediment in still water. Attribute and target based on data from Harvey and Cowx (2003)
Extent and distribution of spawning habitat	m <sup>2</sup> and occurrence	No decline in extent and distribution of spawning beds	Lampreys spawn in clean gravels. Surveys by Inland Fisheries Ireland (IFI) commonly indicated accumulations of redds downstream of major weirs. (See also Gargan et al., 2011)
Availability of juvenile habitat	Number of positive sites in 3rd order channels (and greater), downstream of spawning areas	More than 50% of sample sites positive	Despite observed spawning activity, sampling for ammocoetes consistently fails to find these in many sampling stations and never in any great numbers

Consent of copyright owner required for any other use  
For inspection purposes only

**1096 Brook Lamprey *Lampetra planeri***

**To maintain the favourable conservation condition of Brook Lamprey in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Distribution	% of river accessible	Access to all water courses down to first order streams	Artificial barriers can block or cause difficulties to brook lampreys' migration, both up- and downstream, thereby possibly limiting the species to specific stretches and creating genetically isolated populations (Espanhol et al., 2007)
Population structure of juveniles	Number of age/size groups	At least three age/size groups of brook/river lamprey present	Attribute and target based on data from Harvey and Cowx (2003). It is impossible to distinguish between brook and river lamprey juveniles in the field (Gardiner, 2003), hence they are considered together in this target
Juvenile density in fine sediment	Juveniles/m <sup>2</sup>	Mean catchment juvenile density of brook/river lamprey at least 2/m <sup>2</sup>	Juveniles burrow in areas of fine sediment in still water. Attribute and target based on data from Harvey and Cowx (2003) who state 10/m <sup>2</sup> in optimal conditions and more than 2/m <sup>2</sup> on a catchment basis
Extent and distribution of spawning habitat	m <sup>2</sup> and occurrence	No decline in extent and distribution of spawning beds	Spawning site and redd attributes established by IFI (Rooney et al., in press)
Availability of juvenile habitat	Number of positive sites in 2nd order channels (and greater), downstream of spawning areas	More than 50% of sample sites positive	Many sites with suitable larval attributes i.e. fine sediment in low velocity habitat, are found not to contain larval lamprey. This may be a function of chance or probability, or may be a consequence of insufficient recruitment to fill all spatial niches. Occupancy in excess of 50% of sites would be 'reasonable' for the Irish catchments examined to date (King et al., unpublished data)

1099 River Lamprey *Lampetra fluviatilis*

To maintain the favourable conservation condition of River Lamprey in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	% of river accessible	Access to all water courses down to first order streams	Artificial barriers can block or cause difficulties to river lampreys' migration, both up- and downstream, thereby possibly limiting species to specific stretches and creating genetically isolated populations (Espanhol et al., 2007)
Population structure of juveniles	Number of age/size groups	At least three age/size groups of river/brook lamprey present	Attribute and target based on data from Harvey and Cowx (2003). It is impossible to distinguish between river and brook lamprey juveniles in the field (Gardiner 2003), hence they are considered together in this target
Juvenile density in fine sediment	Juveniles/m <sup>2</sup>	Mean catchment juvenile density of river/brook lamprey at least 2/m <sup>2</sup>	Juveniles burrow in areas of fine sediment in still water. Attribute and target based on data from Harvey and Cowx (2003) who state 10/m <sup>2</sup> in optimal conditions and more than 2/m <sup>2</sup> on a catchment basis
Extent and distribution of spawning habitat	m <sup>2</sup> and occurrence	No decline in extent and distribution of spawning beds	
Availability of juvenile habitat	Number of positive sites in 2nd order channels (and greater), downstream of spawning areas	More than 50% of sample sites positive	Many sites with suitable larval attributes i.e. fine sediment in low velocity habitat, are found not to contain larval lamprey. This may be a function of chance or probability, or may be a consequence of insufficient recruitment to fill all spatial niches. Occupancy in excess of 50% of sites would be 'reasonable' for the Irish catchments examined to date (King et al., unpublished data)

## Conservation objectives for: Lower River Shannon SAC [002165]

### 1106 Atlantic Salmon *Salmo salar* (only in fresh water)

To restore the favourable conservation condition of Salmon in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	% of river accessible	100% of river channels down to second order accessible from estuary	Artificial barriers block salmon's upstream migration, thereby limiting the species to lower stretches and restricting access to spawning areas. The large hydro-electric station at Ardnacrusha and the Parteen regulating weir present considerable obstructions to upstream passage of salmon on the Shannon main channel. While both have fish passes installed, upstream migration of salmon is still problematical. Further weirs upstream on the Shannon also restrict access to spawning habitat. No such obstacles, causing significant fish passage issues for salmon are present on the Feale and Mulkear rivers
Adult spawning fish	Number	Conservation Limit (CL) for each system consistently exceeded	A conservation limit is defined by the North Atlantic Salmon Conservation Organisation (NASCO) as "the spawning stock level that produces long-term average maximum sustainable yield as derived from the adult to adult stock and recruitment relationship". The target is based on the Standing Scientific Committee of the National Salmon Commission's annual model output of CL attainment levels. See SSC (2010). Stock estimates are either derived from direct counts of adults (rod catch, fish counter) or indirectly by fry abundance counts. The salmon stocks in the Shannon above the impoundments are significantly below their Conservation Limits. Salmon stocks in the Feale and Mulkear rivers are above CL
Salmon fry abundance	Number of fry/5 minutes electrofishing	Maintain or exceed 0+ fry mean catchment-wide abundance threshold value. Currently set at 17 salmon fry/5 min sampling	Target is threshold value for rivers currently exceeding their conservation limit (CL). The abundance of salmon fry at monitored sites on the Shannon main channel, above the hydro-electric station, is significantly below this target
Out-migrating smolt abundance	Number	No significant decline	Smolt abundance can be negatively affected by a number of impacts such as estuarine pollution, predation and sea lice ( <i>Lepeophtheirus salmonis</i> ). On the Shannon main channel, salmon smolt abundance may be significantly affected by mortality passing through hydro-electric turbines
Number and distribution of redds	Number and occurrence	No decline in number and distribution of spawning redds due to anthropogenic causes	Salmon spawn in clean gravels. Artificial barriers are currently preventing salmon from accessing suitable spawning habitat on the Shannon main channel

For inspection purposes only.  
Consent of copyright owner required for any other use

**Conservation objectives for: Lower River Shannon SAC [002165]**

**1106 Atlantic Salmon *Salmo salar* (only in fresh water)**

To restore the favourable conservation condition of Salmon in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Water quality	EPA Q value	At least Q4 at all sites sampled by EPA	Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)

For inspection purposes only.  
Consent of copyright owner required for any other use.

**Conservation objectives for: Lower River Shannon SAC [002165]**

**1110 Sandbanks which are slightly covered by sea water all the time**

To maintain the favourable conservation condition of Sandbanks which are slightly covered by sea water all the time in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat distribution	Occurrence	The distribution of sandbanks is stable, subject to natural processes. See map 3	Distribution established using the Valentia Island to River Shannon Admiralty Chart (no. 1819_0)
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 3	Habitat area was estimated as 1,353ha using the Valentia Island to River Shannon Admiralty Chart (no. 1819_0)
Community distribution	Hectares	Conserve the following community type in a natural condition: Subtidal sand to mixed sediment with <i>Nephtys</i> spp. community complex. See map 9	The likely area of the community was derived from a sandbank survey in 2007 (Aquafact, 2007) and a subtidal survey in 2010 (Aquafact, 2011a). See marine supporting document for further details

For inspection purposes only.  
Consent of copyright owner required for any other use.

1130 Estuaries

To maintain the favourable conservation condition of Estuaries in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 4	Habitat area was estimated as 24,273ha using OSi data and the Transitional Water Body area as defined under the Water Framework Directive
Community distribution	Hectares	Conserve the following community types in a natural condition: Intertidal sand to mixed sediment with polychaetes, molluscs and crustaceans community complex; Estuarine subtidal muddy sand to mixed sediment with gammarids community complex; Subtidal sand to mixed sediment with <i>Nucula nucleus</i> community complex; Subtidal sand to mixed sediment with <i>Nephtys</i> spp. community complex; Furoid-dominated intertidal reef community complex; Faunal turf-dominated subtidal reef community; and Anemone-dominated subtidal reef community. See map 9	The likely area of these communities was derived from intertidal and subtidal surveys undertaken in 2010 (Aquafact, 2011a and c). See marine supporting document for further details

For inspection only. Not for publication. Consent of copyright owner published in other use.

**1140 Mudflats and sandflats not covered by seawater at low tide**

To maintain the favourable conservation condition of Mudflats and sandflats not covered by seawater at low tide in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 5	Habitat area was estimated using OSi data as 8,808ha
Community distribution	Hectares	Conserve the following community types in a natural condition: Intertidal sand with <i>Scolelepis squamata</i> and <i>Pontocrates</i> spp. community; and Intertidal sand to mixed sediment with polychaetes, molluscs and crustaceans community complex. See map 9	The likely area of these communities was derived from an intertidal survey in 2010 (Aquafact, 2011c). See marine supporting document for further details

For inspection purposes only.  
Consent of copyright owner required for any other use.



**1150 \*Coastal lagoons**

**To restore the favourable conservation condition of Coastal lagoons in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Habitat area	Hectares	Area stable or increasing, subject to natural processes. Favourable reference area 33.4ha- Shannon Airport Lagoon 24.2ha; Cloonconeen Pool 3.9ha; Scatterry Lagoon 2.8ha; Quayfield and Poulaweala Loughs 2.5ha. See map 6	Areas calculated from spatial data derived from Oliver, 2007. Site codes IL031- IL034. See lagoon supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 6	Sites IL031-IL034 in Oliver, 2007. See lagoon supporting document for further details
Salinity regime	practical salinity units (psu)	Median annual salinity and temporal variation within natural ranges	The lagoons in the site vary from oligohaline to euhaline. See lagoon supporting document for further details
Hydrological regime	Metres	Annual water level fluctuations and minima within natural ranges	Lagoons listed for this site are all considered to be shallow. See lagoon supporting document for further details
Barrier: connectivity between lagoon and sea	Permeability	Appropriate hydrological connections between lagoons and sea, including where necessary, appropriate management	The lagoons within this site exhibit a variety of barrier types including cobble/shingle, karst and artificial embankment. See lagoon supporting document for further details
Water quality: chlorophyll a	µg/L	Annual median chlorophyll a within natural ranges and less than 5µg/L	Target based on Roden and Oliver (2010). See lagoon supporting document for further details
Water quality: Molybdate Reactive Phosphorus (MRP)	mg/L	Annual median MRP within natural ranges and less than 0.1mg/L	Target based on Roden and Oliver (2010). See lagoon supporting document for further details
Water quality: Dissolved Inorganic Nitrogen (DIN)	mg/L	Annual median DIN within natural ranges and less than 0.15mg/L	Target based on Roden and Oliver, 2010). See lagoon supporting document for further details
Depth of macrophyte colonisation	Metres	Macrophyte colonisation to maximum depth of lagoons	As these lagoons are all shallow, it is expected the macrophytes should extend to their deepest points. See lagoon supporting document for further details
Typical plant species	number and m <sup>2</sup>	Maintain number and extent of listed lagoonal specialists, subject to natural variation	Species listed in Oliver, 2007. See lagoon supporting document for further details
Typical animal species	number	Maintain listed lagoon specialists, subject to natural variation	Species listed in Oliver, 2007. See lagoon supporting document for further details
Negative indicator species	Number and % cover	Negative indicator species absent or under control	Low salinity, shallow water and elevated nutrient levels increase the threat of un-natural encroachment by reedbeds

**1160 Large shallow inlets and bays**

To maintain the favourable conservation condition of Large shallow inlets and bays in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 7	Habitat area was estimated as 35,282ha using OSi data and the Transitional Water Body area as defined under the Water Framework Directive
Community distribution	Hectares	Conserve the following community types in a natural condition: Intertidal sand with <i>Scolelepis squamata</i> and <i>Pontocrates</i> spp. community; Intertidal sand to mixed sediment with polychaetes, molluscs and crustaceans community complex; Subtidal sand to mixed sediment with <i>Nucula nucleus</i> community complex; Subtidal sand to mixed sediment with <i>Nephtys</i> spp. community complex; Fucoid-dominated intertidal reef community complex; Mixed subtidal reef community complex; Faunal turf-dominated subtidal reef community; Anemone-dominated subtidal reef community; and <i>Laminaria</i> -dominated community complex. See map 9	The likely area of these communities was derived from intertidal and subtidal surveys in 2010 (Aquafact, 2011a and c). See marine supporting document for further details

For internal use only. Not for public use. Consent of copyright holder is required for any further use.

**1170 Reefs**

To maintain the favourable conservation condition of Reefs in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat distribution	Occurrence	The distribution of Reefs is stable, subject to natural processes. See map 8	Distribution is established from intertidal and subtidal reef surveys in 2010 (Aquafact, 2011b and c)
Habitat area	Hectares	The permanent habitat area is stable, subject to natural processes. See map 8	Habitat area was estimated as 21,421ha from the 2010 intertidal and subtidal reef survey (Aquafact 2011b and c)
Community distribution	Hectares	Conserve the following reef community types in a natural condition: Furoid-dominated intertidal reef community complex; Mixed subtidal reef community complex; Faunal turf-dominated subtidal reef community; Anemone-dominated subtidal reef community; and <i>Laminaria</i> -dominated community complex. See map 9	Based on the 2010 intertidal and subtidal reef survey (Aquafact, 2011b and c). See marine supporting document for further details

For inspection purposes only.  
Consent of copyright owner required for any other use.

**1220 Perennial vegetation of stony banks**

To maintain the favourable conservation condition of Perennial vegetation of stony banks in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession	Current area unknown. It was recorded to be present but extent was not mapped from nine sub-sites during the National Shingle Beach Survey (Moore and Wilson, 1999): Ross Bay, Kilbaha Bay, Clooncneen Lough and Rinevella Bay, Carrigholt Bay, Ballymacrinan Bay, Bunaclugga Bay, Corcas and Sandhills, Bromore and Ballybunnion. NB further unsurveyed areas maybe present within the site
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 10 for recorded locations	Full distribution currently unknown. An excellent array of shingle beaches is known to occur, including three that are ranked of high interest (Ross Bay, Bunaclugga Bay and Clooncneen Lough and Rinevella), the last of which is associated with a lagoonal system (Moore and Wilson, 1999). Habitat likely to be more widespread. See coastal habitats supporting document for further details. See also the conservation objective for coastal lagoons (1150)
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Based on data from Moore and Wilson (1999). Shingle features are relatively stable in the long-term and shingle beaches within this SAC appear to be functioning naturally with few artificial restrictions to beach dynamics (Moore and Wilson, 1999). See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Moore and Wilson (1999). Lichens are present at Ross Bay and Clooncneen and Rinevella Bay indicating a degree of stability. See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain the typical vegetated shingle flora including the range of sub-communities within the different zones	The Carrigaholt sub-site is a small site with a diverse flora. The Bunaclugga Bay sub-site supports yellow horned-poppy ( <i>Glaucium flavum</i> ), which contributes to the site's high interest ranking. Based on data from Moore and Wilson (1999). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Moore and Wilson (1999). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. See coastal habitats supporting document for further details

**Conservation objectives for: Lower River Shannon SAC [002165]**

**1230 Vegetated sea cliffs of the Atlantic and Baltic coasts**

To maintain the favourable conservation condition of Vegetated sea cliffs in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat length	Kilometres	Area stable or increasing, subject to natural processes, including erosion. For sub-sites mapped: Kilbaha- 4.1km; Ladder Rock- 1.0km; Moyarta- 0.9km; Lisheencrony- 1.1km; Burrane- 0.2km; Kerry Head- 33.4km; Ballybunion- 15.6km; Kilclogher- 4.9km; Loop Head- 6.1km. See map 11	Based on data from the Irish Sea Cliff Survey (ISCS) (Barron et al., 2011). Nine sub-sites were identified using a combination of aerial photos and the DCENR helicopter viewer. The length of each cliff was measured (in some cases the cliff was measured in sections) to give a total estimated area of 67.3km within the SAC. Cliffs are linear features and are therefore measured in kilometres. Length of cliff likely to be underestimated. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 11	Based on data from the Irish Sea Cliff Survey (Barron et al., 2011). Most of the SAC west of Kilcredaun Point and Kilconly Point is bounded by high rocky sea cliffs. Both hard and soft cliffs occur in this SAC (ISCS; Browne, 2005). See coastal habitats supporting document for further details
Physical structure: functionality and hydrological regime	Occurrence of artificial barriers	No alteration to natural functioning of geomorphological and hydrological processes due to artificial structures	Based on data from the Irish Sea Cliff Survey (Barron et al., 2011). Maintaining natural geomorphological processes including natural erosion is important for the health of vegetated sea cliff. Hydrological processes maintain flushes and in some cases tufa formations that can be associated with sea cliffs. Freshwater seepage was noted from the cliffs at Loop Head and Kilclogher. Stream or cascade was noted from Kerry Head. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain range of sea cliff habitat zonation including transitional zones, subject to natural processes including erosion and succession	Based on data from the Irish Sea Cliff Survey (Barron et al., 2011). At Loop Head sub-site the zones recorded were: splash, crevice ledge and ungrazed coastal grassland on hard cliffs. At Kerry Head sub-site the zones recorded were: splash, pioneer, crevice ledge, ungrazed/grazed coastal grassland on hard cliffs and coastal grassland on soft cliffs. See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from the Irish Sea Cliff Survey (Barron et al., 2011). See coastal habitats supporting document for further details

**1230 Vegetated sea cliffs of the Atlantic and Baltic coasts**

To maintain the favourable conservation condition of Vegetated sea cliffs in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in the Irish Sea cliff survey (Barron et al., 2011)	Based on data from the Irish Sea Cliff Survey (Barron et al., 2011). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from the Irish Sea Cliff Survey (Barron et al., 2011). See coastal habitats supporting document for further details
Vegetation composition: bracken and woody species	Percentage	Cover of bracken ( <i>Pteridium aquilinum</i> ) on grassland and/or heath to be less than 10%. Cover of woody species on grassland and/or heath to be less than 20%	Based on data from the Irish Sea Cliff Survey (Barron et al., 2011). See coastal habitats supporting document for further details

For inspection purposes only.  
Consent of copyright owner required for any other use.

**Conservation objectives for: Lower River Shannon SAC [002165]**

**1310 *Salicornia* and other annuals colonizing mud and sand**

To maintain the favourable conservation condition of *Salicornia* and other annuals colonizing mud and sand in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Carrigafoyle - 0.005ha; Inishdea, Owenshere - 0.003ha; Knock - 0.029ha; Querin - 0.185ha; Rinevilla Bay - 0.001ha. See map 12	Based on data from Saltmarsh Monitoring Project (SMP) (McCorry and Ryle, 2009). Habitat recorded at five of the ten sub-sites surveyed and mapped, giving a total estimated area of 0.223ha. NB further unsurveyed areas may be present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 12 for known distribution	Based on data from McCorry and Ryle (2009). Habitat recorded at six out of ten sub-sites by McCorry and Ryle (2009). NB further unsurveyed areas may be present within the site. <i>Salicornia</i> is an annual species, so its distribution can vary significantly from year to year. See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/ absence of physical barriers	Maintain natural circulation of sediments and organic matter without any physical obstructions	Sediment supply is particularly important for this pioneer saltmarsh community, as the distribution of this habitat depends on accretion rates. See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain/restore creek and pan structure, subject to natural processes, including erosion and succession	Based on data from McCorry and Ryle (2009). Creeks deliver sediment throughout saltmarsh system. Creeks and pan structures well developed in the larger sections of the marsh at Carrigafoyle, Shepperton/Fergus Estuary and Inishdea/Owenshere. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	This pioneer saltmarsh community requires regular tidal inundation. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimeters	Maintain structural variation within sward	Based on data from McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% of area outside creeks vegetated	Based on data from McCorry and Ryle (2009). See coastal habitats supporting document for further details

**Conservation objectives for: Lower River Shannon SAC [002165]**

**1310 *Salicornia* and other annuals colonizing mud and sand**

To maintain the favourable conservation condition of *Salicornia* and other annuals colonizing mud and sand in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation composition: typical species and sub-communities	Percentage cover	Maintain the presence of species-poor communities with typical species listed in Saltmarsh Monitoring Project (McCorry and Ryle, 2009)	Based on data from McCorry and Ryle (2009). Species of local distinctiveness recorded include sea wormwood ( <i>Seriphidium maritimum</i> ), meadow barley ( <i>Hordeum secalinum</i> ) and hard grass ( <i>Parapholis strigosa</i> ) (McCorry and Ryle, 2009; internal NPWS files). See coastal habitats supporting document for further details
Vegetation structure: negative indicator species- <i>Spartina anglica</i>	Hectares	No significant expansion of common cordgrass ( <i>Spartina anglica</i> ), with an annual spread of less than 1%	Based on data from McCorry and Ryle (2009). <i>Spartina</i> was recorded at all sub-sites and is considered a significant threat to the habitat. See coastal habitats supporting document for further details

For inspection purposes only.  
Consent of copyright owner required for any other use.



**Conservation objectives for: Lower River Shannon SAC [002165]**

**1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)**

To restore the favourable conservation condition of Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Carrigafoyle- 6.774ha; Barrigone, Aughinish- 10.288ha; Beagh- 0.517ha; Bunratty- 26.939ha; Shepperton, Fergus Estuary- 37.925ha; Inishdea, Owenshere- 18.127ha; Killadysert, Inishcorker- 2.604ha; Knock- 0.576ha; Querin- 3.726ha; Rinevilla Bay- 11.883ha. See map 12	Based on data from the Saltmarsh Monitoring Project (SMP) (McCorry and Ryle 2009). Ten sub-sites that supported Atlantic salt meadow were mapped (119.36ha) and additional areas of potential saltmarsh (376.07ha) were identified from an examination of aerial photographs, giving a total estimated area of 495.43ha. Saltmarsh habitat also occurs at 11 other sub-sites within the SAC (Curtis and Sheehy-Skeffington, 1998). NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 12 for mapped distribution	Based on data from McCorry and Ryle (2009). Within the sites surveyed by the SMP, estuary type saltmarsh over a mud substrate is most common and ASM is the dominant saltmarsh habitat. See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/ absence of physical barriers	Maintain natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry and Ryle (2009). Embankments along much of the shoreline are a feature of this SAC. These embankments were erected in the past and much of the site has been remodelled and large areas of land reclaimed as a result. See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from McCorry and Ryle (2009). Creeks and pan structures well developed at the larger sections of ASM in the Carrigafoyle sub-site. At the ASM at Shepperton, Fergus Estuary, the larger patches still retain a natural creek and salt pan structure. At Inishdea, Owenshere sub-site within some of the intact saltmarsh, there is a complex network of creeks, salt pans and depressions. At Killadysart, Inishcorker and Querin, creek and pan development is generally poor. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	See coastal habitats supporting document for further details

**Conservation objectives for: Lower River Shannon SAC [002165]**

**1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)**

To restore the favourable conservation condition of Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry and Ryle (2009). Zonations to other saltmarsh habitats as well as brackish and terrestrial habitats were recorded at all sub-sites. See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimeters	Maintain structural variation within sward	Based on data from McCorry and Ryle (2009). All of the sub-sites are grazed to some extent. Overgrazing was noted from Carrigafoyle, Shepperton, Fergus Estuary and Knock sub-sites. See coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% of the saltmarsh area vegetated	Based on data from McCorry and Ryle (2009). Some poaching was noted from most of the sub-sites. See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in Saltmarsh Monitoring Project (McCorry and Ryle 2009)	See coastal habitats supporting document for further details
Vegetation structure: negative indicator species- <i>Spartina anglica</i>	Hectares	No significant expansion of common cordgrass ( <i>Spartina anglica</i> ), with an annual spread of less than 1%	Based on data from McCorry and Ryle (2009). <i>Spartina</i> is a major element of the vegetation at all sub-sites in this SAC. See coastal habitats supporting document for further details

1349 Bottlenose Dolphin *Tursiops truncatus*

To maintain the favourable conservation condition of Bottlenose Dolphin in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Access to suitable habitat	Number of artificial barriers	Species range within the site should not be restricted by artificial barriers to site use. See map 16 for suitable habitat	See marine supporting document for further details
Habitat use: critical areas	Location and hectares	Critical areas, representing habitat used preferentially by bottlenose dolphin, should be maintained in a natural condition. See map 16	Attribute and target based on Ingram and Rogan (2002), Englund et al. (2007), Englund et al. (2008), Berrow (2009), Berrow et al. (2010) and review of data from other studies. See marine supporting document for further details
Disturbance	Level of impact	Human activities should occur at levels that do not adversely affect the bottlenose dolphin population at the site	See marine supporting document for further details

For inspection purposes only.  
Consent of copyright owner required for any other use.

**1355 Otter *Lutra lutra***

**To restore the favourable conservation condition of Otter in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. FCS target, based on 1980/81 survey findings, is 88% in SACs. Current range in Shannon catchment estimated at 70.5% (Bailey and Rochford 2006)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 596.8ha above high water mark (HWM); 958.9ha along river banks/ around ponds	No field survey. Areas mapped to include 10m terrestrial buffer along shoreline (above HWM and along river banks) identified as critical for otters (NPWS, 2007)
Extent of marine habitat	Hectares	No significant decline. Area mapped and calculated as 4,461.6ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (HWM) (NPWS, 2007; Kruuk, 2006)
Extent of freshwater (river) habitat	Kilometers	No significant decline. Length mapped and calculated as 500.1km	No field survey. River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake/lagoon) habitat	Hectares	No significant decline. Area mapped and calculated as 125.6ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk, 2006; Kruuk and Moorhouse, 1991)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006) and wrasse and rockling in coastal waters (Kingston et al., 1999)
Barriers to connectivity	Number	No significant increase. For guidance, see map 17	Otters will regularly commute across stretches of open water up to 500m. e.g. between the mainland and an island; between two islands; across an estuary (De Jongh and O'Neill, 2010). It is important that such commuting routes are not obstructed

**Conservation objectives for: Lower River Shannon SAC [002165]**

**1410 Mediterranean salt meadows (*Juncetalia maritimi*)**

To restore the favourable conservation condition of Mediterranean salt meadows (*Juncetalia maritimi*) in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Carrigafoyle- 4.193ha; Barrigone, Aghinish- 2.407ha; Bunratty- 0.865ha; Inishdea, Owenshere- 11.609ha; Killadysert, Inishcorker- 0.705ha; Knock- 0.143ha, Querin- 0.008ha; Rinevilla Bay- 2.449ha. See map 12	Based on data from the Saltmarsh Monitoring Project (SMP) (McCorry and Ryle, 2009). Eight sub-sites that support Mediterranean salt meadow were mapped (22.379ha) and additional areas of potential saltmarsh (25.646ha) were identified from an examination of aerial photographs, giving a total estimated area of 48.025ha. Saltmarsh habitat also occurs at 11 other sub-sites within the SAC (Curtis and Sheehy-Skeffington, 1998). NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 12 for known distribution	Based on data from McCorry and Ryle (2009). Within the sites surveyed by the SMP, estuary type saltmarsh over a mud substrate is most common. See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain natural circulation of sediments and organic matter, without any physical obstructions	Based on data from McCorry and Ryle (2009). Embankments along much of the shoreline are a feature of this SAC. These embankments were erected in the past and much of the site has been remodelled and large areas of land reclaimed because of them. See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain/restore creek and pan structure, subject to natural processes, including erosion and succession	Based on data from the Saltmarsh Monitoring Project (McCorry and Ryle, 2009). The MSM at Carrigafoyle contains some large salt pans. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	Mediterranean salt meadow is found high up in the saltmarsh but requires occasional tidal inundation. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry and Ryle (2009). Zonations to other saltmarsh habitats as well as brackish and terrestrial habitats were recorded at most sub-sites. See coastal habitats supporting document for further details

**Conservation objectives for: Lower River Shannon SAC [002165]**

**1410 Mediterranean salt meadows (*Juncetalia maritimi*)**

To restore the favourable conservation condition of Mediterranean salt meadows (*Juncetalia maritimi*) in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from McCorry and Ryle (2009). All of the sub-sites are grazed to some extent. Overgrazing was noted from Inishdea, Owenshere and Knock sub-sites. See coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% of area outside creeks vegetated	Based on data from McCorry and Ryle (2009). Some poaching was noted from most of the sub-sites. See coastal habitats supporting document for further details
Vegetation composition: typical species	Percentage cover	Maintain range of sub-communities with typical species listed in Saltmarsh Monitoring Project (McCorry and Ryle, 2009)	See coastal habitats supporting document for further details
Vegetation structure: negative indicator species - <i>Spartina anglica</i>	Hectares	No significant expansion of common cordgrass ( <i>Spartina anglica</i> ), with an annual spread of less than 1%	Based on data from McCorry and Ryle (2009). <i>Spartina</i> is a major element of the vegetation at all sub-sites in this SAC. See coastal habitats supporting document for further details

For inspection purposes only. Consent of copyright owner required for any other use.

**Conservation objectives for: Lower River Shannon SAC [002165]**

**3260 Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation**

To maintain the favourable conservation condition of Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Kilometres	Area stable or increasing, subject to natural processes	Three sub-types of high conservation value are known to occur in the site. See Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation supporting document for further details. <b>Note: rooted macrophytes should be absent or trace (&lt; 5% cover) in freshwater pearl mussel (<i>Margaritifera margaritifera</i>) habitat. The freshwater pearl mussel (1029) conservation objective takes precedence over this objective for habitat 3260 in the Cloon River within this SAC, because the mussel requires environmental conditions closer to natural background levels</b>
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 13	See Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation supporting document for further details
Hydrological regime: river flow	Metres per second	Maintain appropriate hydrological regimes	See Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation supporting document for further details
Hydrological regime: tidal influence	Daily water level fluctuations - metres	Maintain natural tidal regime	See Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation supporting document for further details
Hydrological regime: freshwater seepages	Metres per second	Maintain appropriate freshwater seepage regimes	See Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation supporting document for further details
Substratum composition: particle size range	Millimetres	The substratum should be dominated by the particle size ranges, appropriate to the habitat sub-type (frequently sands, gravels and cobbles)	Although many of the high-conservation-value sub-types are dominated by coarse substrata, for certain sub-types, notably triangular club-rush ( <i>Schoenoplectus triqueter</i> ) and opposite-leaved pondweed ( <i>Groenlandia densa</i> ), fine substrata are required. See Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation supporting document for further details

**Conservation objectives for: Lower River Shannon SAC [002165]**

**3260 Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation**

To maintain the favourable conservation condition of Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Water quality: nutrients	Milligrammes per litre	The concentration of nutrients in the water column should be sufficiently low to prevent changes in species composition or habitat condition	The specific targets may vary among sub-types. See Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation supporting document for further details
Vegetation composition: typical species	Occurrence	Typical species of the relevant habitat sub-type should be present and in good condition	See Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation supporting document for further details
Floodplain connectivity	Area	The area of active floodplain at and upstream of the habitat should be maintained	See Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation supporting document for further details
Riparian habitat	Area	The area of riparian woodland at and upstream of the bryophyte-rich sub-type should be maintained	See Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation supporting document for further details. See also the conservation objective for Alluvial forests with <i>Alnus glutinosa</i> and <i>Fraxinus excelsior</i> ( <i>Alno-Padion</i> , <i>Alnion incanae</i> , <i>Salicion albae</i> ) (91E0)

For inspection purposes only. Consent of copyright owner required for any other use.



**Conservation objectives for: Lower River Shannon SAC [002165]**

**6410 *Molinia* meadows on calcareous, peaty or clayey-silt-laden soils (*Molinion caeruleae*)**

To maintain the favourable conservation condition of *Molinia* meadows on calcareous, peaty or clayey-silt laden soils (*Molinion caeruleae*) in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Full extent of this habitat in this site is currently unknown- see distribution below
Habitat distribution	Occurrence	No decline, subject to natural processes	This habitat has been recorded on the eastern bank of the Shannon, just north of Castleconnell, Co. Limerick (NPWS internal files). Full distribution of this habitat in this site is currently unknown and it almost certainly occurs elsewhere. The Irish semi-natural grasslands survey will cover Co. Limerick in 2012 and additional information is likely to be available following this survey
Vegetation structure: broadleaf herb: grass ratio	Percentage	Broadleaf herb component of vegetation between 40 and 90%	Attribute and target based on O'Neill et al. (2010)
Vegetation structure: sward height	Percentage	30-70% of sward between 10 and 80cm high	Attribute and target based on O'Neill et al. (2010)
Vegetation composition: typical species	Number	At least 7 positive indicator species present, including 1 "high quality" species	List of positive indicator species, including high quality species, identified by O'Neill et al. (2010). Note that purple moor-grass ( <i>Molinia caerulea</i> ) is a positive indicator species, but not necessarily an essential component of the habitat
Vegetation composition: notable species	Number	No decline, subject to natural processes	A number of notable species have been recorded in this habitat at this site including smooth brome ( <i>Bromus racemosus</i> ), pale sedge ( <i>Carex pallescens</i> ) and blue-eyed grass ( <i>Sisyrinchium bermudiana</i> ) (Reynolds et al., 2006)
Vegetation composition: negative indicator species	Percentage	Negative indicator species collectively not more than 20% cover, with cover by an individual species less than 10%. Non-native invasive species, absent or under control	List of negative indicator species identified by O'Neill et al. (2010)
Vegetation composition: negative indicator moss species	Percentage	Bog mosses ( <i>Sphagnum</i> spp.) not more than 10% cover; hair mosses ( <i>Polytrichum</i> spp.) not more than 25% cover	Attribute and target based on O'Neill et al. (2010)

**Conservation objectives for: Lower River Shannon SAC [002165]**

**6410 *Molinia* meadows on calcareous, peaty or clayey-silt-laden soils (*Molinion caeruleae*)**

To maintain the favourable conservation condition of *Molinia* meadows on calcareous, peaty or clayey-silt laden soils (*Molinion caeruleae*) in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation structure: woody species and bracken ( <i>Pteridium aquilinum</i> )	Percentage	Cover of woody species and bracken not more than 5% cover	Attribute and target based on O'Neill et al. (2010)
Physical structure: bare ground	Percentage	Not more than 10% bare ground	Attribute and target based on O'Neill et al. (2010)

For inspection purposes only.  
Consent of copyright owner required for any other use.

**Conservation objectives for: Lower River Shannon SAC [002165]**

**91E0 \*Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (*Alno-Padion*, *Alnion incanae*, *Salicion albae*)**

To restore the favourable conservation condition of Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (*Alno-Padion*, *Alnion incanae*, *Salicion albae*) in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, at least c.8.5ha for sites surveyed. See map 14	Minimum area, based on 5 sites surveyed by Perrin et al. (2008) - site codes 1286, 1577, 1857, 1861, 1995. See woodland habitats supporting document for further details. NB further areas are likely to be present within the SAC
Habitat distribution	Occurrence	No decline. Surveyed locations shown on map 14	Distribution based on Perrin et al. (2008). NB further areas are likely to be present within the SAC
Woodland size	Hectares	Area stable or increasing. Where topographically possible, "large" woods at least 25ha in size and "small" woods at least 3ha in size	The sizes of at least some of the existing woodlands need to be increased in order to reduce habitat fragmentation and benefit those species requiring 'deep' woodland conditions (Peterken, 2002). Topographical and land-ownership constraints may restrict expansion
Woodland structure: cover and height	Percentage and metres	Diverse structure with a relatively closed canopy containing mature trees; subcanopy layer with semi-mature trees and shrubs; and well-developed herb layer	Described in Perrin et al. (2008). See woodland habitats supporting document for further details
Woodland structure: community diversity and extent	Hectares	Maintain diversity and extent of community types	Described in Perrin et al. (2008). See woodland habitats supporting document for further details
Woodland structure: natural regeneration	Seedling: sapling: pole ratio	Seedlings, saplings and pole age-classes occur in adequate proportions to ensure survival of woodland canopy	Alder and oak regenerate poorly. Ash often regenerates in large numbers although few seedlings reach pole size
Hydrological regime: flooding depth/height of water table	Metres	Appropriate hydrological regime necessary for maintenance of alluvial vegetation	Periodic flooding is essential to maintain alluvial woodlands along river floodplains
Woodland structure: dead wood	m <sup>3</sup> per hectare; number per hectare	At least 30m <sup>3</sup> /ha of fallen timber greater than 10cm diameter; 30 snags/ha; both categories should include stems greater than 40cm diameter (greater than 20cm diameter in the case of alder)	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem

**Conservation objectives for: Lower River Shannon SAC [002165]**

**91E0 \*Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (Alno-Padion, *Alnion incanae*, *Salicion albae*)**

To restore the favourable conservation condition of Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (Alno-Padion, *Alnion incanae*, *Salicion albae*) in the Lower River Shannon SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Woodland structure: veteran trees	Number per hectare	No decline	Mature and veteran trees are important habitats for bryophytes, lichens, saproxylic organisms and some bird species. Their retention is important to ensure continuity of habitats/niches and propagule sources
Woodland structure: indicators of local distinctiveness	Occurrence	No decline	Includes ancient or long-established woodlands, archaeological and geological features as well as red-data and other rare or localised species. Perrin and Daly (2010) list four sites as containing potential ancient/long established woodland. See woodland habitats supporting document for further details
Vegetation composition: native tree cover	Percentage	No decline. Native tree cover not less than 95%	Species reported in Perrin et al. (2008)
Vegetation composition: typical species	Occurrence	A variety of typical native species present, depending on woodland type, including alder ( <i>Alnus glutinosa</i> ), willows ( <i>Salix</i> spp) and, locally, oak ( <i>Quercus robur</i> ) and ash ( <i>Fraxinus excelsior</i> )	Species reported in Perrin et al. (2008). See woodland habitats supporting document for further details
Vegetation composition: negative indicator species	Occurrence	Negative indicator species, particularly non-native invasive species, absent or under control	The following are the most common invasive species in this woodland type: Himalayan balsam ( <i>Impatiens glandulifera</i> ), giant hogweed ( <i>Heracleum mantegazzianum</i> ), sycamore ( <i>Acer pseudoplatanus</i> )



**Legend**  
 SAC 002165

An Roinn Ealaíon, Oidhreacht agus Gaeltachta  
 Department of Arts, Heritage and the Gaeltacht


**MAP 1:  
 LOWER RIVER SHANNON SAC  
 CONSERVATION OBJECTIVES  
 SAC DESIGNATION**  
 Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE: SAC 002165  
 CO. CLARE; version 1.2, CO. CORK; version 1.01, CO. KERRY; version 1.11,  
 CO. LIMERICK; version 1.11, CO. TIPPERARY; version 1.05

0 5 10 15 20 km


The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).  
 Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaite. Macasamhail d'ábhar na Suirbhéarachtá Ordoináis le chead ón Rialtas (Ceadúnas Uimh. EN 0059208)


**Map Version 1**  
 Date: June 2012



**Legend**


- 002165 Lower River Shannon SAC
- 000930 Clare Glen SAC
- 004077 River Shannon and River Fergus Estuaries SPA
- 004119 Loop Head SPA
- 004161 Stack's to Mullaghareirk Mountains, West Limerick Hills and Mount Eagle SPA
- 004165 Slievefelim to Silvermines Mountains SPA
- 004189 Kerry Head SPA
- OSi Discovery Series County Boundaries





**Legend**

- SAC 002165
- 1130 Estuaries
- OSi Discovery Series County Boundaries


 **An Roinn Ealaíon, Oidhreacht agus Gaeltachta**  
 Department of Arts, Heritage and the Gaeltacht

**MAP 4:  
 LOWER RIVER SHANNON SAC  
 CONSERVATION OBJECTIVES  
 ESTUARIES**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE: SAC 002165  
 CO. CLARE; version 1.2, CO. CORK; version 1.01, CO. KERRY; version 1.11,  
 CO. LIMERICK; version 1.11, CO. TIPPERARY; version 1.05


0    2    4    6    8    10 km



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).  
 Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Macasamhail d'ábhar na Suirbhéarachta Ordonáis le chead ón Rialtas (Ceadúnas Uimh. EN 0059208)


  
**Map Version 1**  
**Date: June 2012**





**Legend**

- SAC 002165
- 1140 Mudflats and sandflats not covered by sea water at low tide
- OSi Discovery Series County Boundaries

 **An Roinn Ealaíon, Oidhreacht agus Gaeltachta**  
 Department of Arts, Heritage and the Gaeltacht

**MAP 5:**  
**LOWER RIVER SHANNON SAC**  
**CONSERVATION OBJECTIVES**  
**TIDAL MUDFLATS AND SANDFLATS**


Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE: SAC 002165  
 CO. CLARE; version 1.2, CO. CORK; version 1.01, CO. KERRY; version 1.11,  
 CO. LIMERICK; version 1.11, CO. TIPPERARY; version 1.05

0      5      10      15 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).  
 Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Macasamhail d'ábhar na Suirbhéarachta Ordonáis le chead ón Rialtas (Ceadúnas Uimh. EN 0059208)


  
**Map Version 1**  
**Date: June 2012**






**Legend**

- SAC 002165
- 1160 Large shallow inlets and bays
- OSi Discovery Series County Boundaries





**Legend**

- SAC 002165
- 1170 Reefs
- OSi Discovery Series County Boundaries

**An Roinn Ealaíon, Oidhreacht agus Gaeltachta**  
 Department of Arts, Heritage and the Gaeltacht

**MAP 8:  
 LOWER RIVER SHANNON SAC  
 CONSERVATION OBJECTIVES  
 REEFS**


Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE: SAC 002165  
 CO. CLARE; version 1.2, CO. CORK; version 1.01, CO. KERRY; version 1.11,  
 CO. LIMERICK; version 1.11, CO. TIPPERARY; version 1.05

0      5      10      15 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).  
 Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Macasamhail d'ábhar na Suirbhéarachta Ordonáis le chead ón Rialtas (Ceadunas Uimh. EN 0059208)

N  
  
**Map Version 1**  
**Date: June 2012**





For inspection purposes only. Consent of copyright owners required for any other use.


**Legend**


- SAC 002165
- OSi Discovery Series County Boundaries

**Marine Community Types**


- Anemone-dominated subtidal reef community
- Estuarine subtidal muddy sand to mixed sediment with gammarids community complex
- Faunal turf-dominated subtidal reef community
- Fucoid-dominated intertidal reef community complex
- Intertidal sand to mixed sediment with polychaetes, molluscs and crustaceans community complex
- Intertidal sand with *Scolelepis squamata* and *Pontocrates* spp. community
- Laminaria*-dominated community complex
- Mixed subtidal reef community complex
- River
- Subtidal sand to mixed sediment with *Nephtys* spp. community complex
- Subtidal sand to mixed sediment with *Nucula nucleus* community complex
















For inspection purposes only.  
Consent of copyright owner required for any other use.

**Legend**

- SAC 002165
- OSi Discovery Series County Boundaries
- NSNW: 1857 National Survey of Native Woodland Site Codes


**Woodland Habitats**

- 91E0 \*Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (Alno-padion, Alnion incanae, Salicion albae) Qualifying Interest
- 91A0 / 91E0 Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles / \*Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (Alno-padion, Alnion incanae, Salicion albae) Qualifying Interest
- 91A0 Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles
- WN2 Oak-ash-hazel woodland




**Legend**

- SAC 002165
- 1029 Freshwater Pearl Mussel - *Margaritifera margaritifera* Suitable Habitat
- 1029 Freshwater Pearl Mussel - *Margaritifera margaritifera* Distribution
- 1029 Freshwater Pearl Mussel - *Margaritifera margaritifera* Catchment
- OSi Discovery Series County Boundaries




**Legend**

- SAC 002165
- 1349 Bottlenose Dolphin - *Tursiops truncatus* Critical Habitat
- 1349 Bottlenose Dolphin - *Tursiops truncatus* Habitat
- OSi Discovery Series County Boundaries



**Legend**

- SAC 002165
- 1355 Otter - *Lutra lutra* 250m Commuting Buffer
- OSI Discovery Series County Boundaries

 **An Roinn Ealaíon, Oidhreacht agus Gaeltachta**  
Department of Arts, Heritage and the Gaeltacht

**MAP 17:**  
**LOWER RIVER SHANNON SAC**  
**CONSERVATION OBJECTIVES**  
**OTTER COMMUTING**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE: SAC 002165  
CO. CLARE; version 1.2, CO. CORK; version 1.01, CO. KERRY; version 1.11,  
CO. LIMERICK; version 1.11, CO. TIPPERARY; version 1.05

0      5      10      15 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).  
Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Macasamhail d'ábhar na Suirbhéarachta Ordonáis le chead ón Rialtas (Ceadunas Uimh. EN 0059208)

  
**Map Version 1**  
**Date: June 2012**

# National Parks and Wildlife Service

## Conservation Objectives Series

### River Shannon and River Fergus Estuaries SPA 004077

For inspection purposes only.  
Consent of copyright owner required for any other use.



**An Roinn**  
**Ealaíon, Oidhreacht agus Gaeltachta**  
**Department of**  
**Arts, Heritage and the Gaeltacht**



**National Parks and Wildlife Service,  
Department of Arts, Heritage and the Gaeltacht,  
7 Ely Place, Dublin 2, Ireland.  
Web: [www.npws.ie](http://www.npws.ie)  
E-mail: [nature.conservation@ahg.gov.ie](mailto:nature.conservation@ahg.gov.ie)**

**Citation:**

NPWS (2012) Conservation Objectives: River Shannon and River Fergus Estuaries SPA 004077. Version 1.0.  
National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht.

**Series Editors: Rebecca Jeffrey & Naomi Kingston  
ISSN 2009-4086**

## Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

### Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.



## Qualifying Interests

\* indicates a priority habitat under the Habitats Directive

### 004077 River Shannon and River Fergus Estuaries SPA

A017	Cormorant <i>Phalacrocorax carbo</i>	breeding + wintering
A038	Whooper Swan <i>Cygnus cygnus</i>	wintering
A046	Light-bellied Brent Goose <i>Branta bernicla hrota</i>	wintering
A048	Shelduck <i>Tadorna tadorna</i>	wintering
A050	Wigeon <i>Anas penelope</i>	wintering
A052	Teal <i>Anas crecca</i>	wintering
A054	Pintail <i>Anas acuta</i>	wintering
A056	Shoveler <i>Anas clypeata</i>	wintering
A062	Scaup <i>Aythya marila</i>	wintering
A137	Ringed Plover <i>Charadrius hiaticula</i>	wintering
A140	Golden Plover <i>Pluvialis apricaria</i>	wintering
A141	Grey Plover <i>Pluvialis squatarola</i>	wintering
A142	Lapwing <i>Vanellus vanellus</i>	wintering
A143	Knot <i>Calidris canutus</i>	wintering
A149	Dunlin <i>Calidris alpina</i>	wintering
A156	Black-tailed Godwit <i>Limosa limosa</i>	wintering
A157	Bar-tailed Godwit <i>Limosa lapponica</i>	wintering
A160	Curlew <i>Numenius arquata</i>	wintering
A162	Redshank <i>Tringa totanus</i>	wintering
A164	Greenshank <i>Tringa nebularia</i>	wintering
A179	Black-headed Gull <i>Chroicocephalus ridibundus</i>	wintering
A999	Wetlands	

**Please note that this SPA overlaps with Lower River Shannon SAC (002165). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping site as appropriate.**

## Supporting documents, relevant reports & publications (listed by date)

Supporting documents, NPWS reports and publications are available for download from: [www.npws.ie/Publications](http://www.npws.ie/Publications)

**Title:** BirdLife International Seabird Ecology and Foraging Range Database

**Year:** 2012

**Author:** BirdLife International

**Series:** <http://seabird.wikispaces.com>

**Title:** Seabird Monitoring Programme (SMP) Database

**Year:** 2012

**Author:** JNCC

**Series:** <http://jncc.defra.gov.uk/smp/Default.aspx>

**Title:** River Shannon and River Fergus Estuaries SPA (004077). Conservation objectives supporting document. [Version 1]

**Year:** 2012

**Author:** NPWS

**Series:** Unpublished Report to NPWS

**Title:** Seabird Populations of Britain and Ireland

**Year:** 2004

**Author:** Mitchell, P.I.; Newton, S.F.; Ratcliffe, N.; Dunn, T.E.

**Series:** Poyser, London

**Title:** Seabird monitoring handbook for Britain and Ireland: a compilation of methods for survey and monitoring of breeding seabirds.

**Year:** 1995

**Author:** Walsh, P.; Halley, D.J.; Harris, M.P.; del Nevo, A.; Sim, I.M.W.; Tasker, M.L.

**Series:** JNCC, Peterborough

For inspection purposes only.  
Consent of copyright owner required for any other use.

**Conservation objectives for: River Shannon and River Fergus Estuaries SPA [004077]**

**A017 Cormorant *Phalacrocorax carbo***

To maintain the favourable conservation condition of Cormorant in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Breeding population abundance: apparently occupied nests (AONs)	Number	No significant decline	This attribute applies to breeding cormorant. Measure based on standard survey methods (see Walsh et al., 1995). Mitchell et al. (2004) provides summary population information. The Seabird Monitoring Programme (SMP) online database (JNCC, 2012) provides population data for this species
Productivity rate	Mean number	No significant decline	This attribute applies to breeding cormorant. Measure based on standard survey methods (see Walsh et al., 1995). The Seabird Monitoring Programme (SMP) online database (JNCC, 2012) provides population data for this species
Distribution: breeding colonies	Number; location; area (hectares)	No significant decline	This attribute applies to breeding cormorant. Cormorant colonies are usually sited on flat or rocky islets or sea stack tops, less often on cliffs but they can also nest in trees (Walsh et al., 1995)
Prey biomass available	Kilogrammes	No significant decline	This attribute applies to breeding cormorant. Key prey items: fish (mostly benthic), some crustaceans. Key habitats: populations use sandy areas, rocky and vegetated substrate. Foraging range: max. 50km, mean max. 31.67km, mean 8.46km (BirdLife International Seabird Database (Birdlife International, 2012))
Barriers to connectivity	Number; location; shape; area (hectares)	No significant increase	This attribute applies to breeding cormorant. Seabird species can make extensive use of the marine waters adjacent to their breeding colonies. Foraging range: max. 50km, mean max. 31.67km, mean 8.46km (BirdLife International Seabird Database (Birdlife International, 2012))
Disturbance at the breeding site	Level of impact	Human activities should occur at levels that do not adversely affect the breeding cormorant population	This attribute applies to breeding cormorant. Cormorant colonies are usually sited on flat or rocky islets or sea stack tops, less often on cliffs but they can also nest in trees (Walsh et al., 1995)
Population trend	Percentage change	Long term population trend stable or increasing	This attribute applies to non-breeding cormorant. Waterbird population trends are presented in part four of the conservation objectives supporting document

**A017 Cormorant *Phalacrocorax carbo***

**To maintain the favourable conservation condition of Cormorant in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by cormorant other than that occurring from natural patterns of variation	This attribute applies to non-breeding cormorant. As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A038 Whooper Swan *Cygnus cygnus***

**To maintain the favourable conservation condition of Whooper Swan in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by whooper swan other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A046 Light-bellied Brent Goose *Branta bernicla hrota***

To maintain the favourable conservation condition of Light-bellied Brent Goose in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by light-bellied brent goose other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

For inspection purposes only.  
Consent of copyright owner required for any other use.

**A048 Shelduck *Tadorna tadorna***

**To maintain the favourable conservation condition of Shelduck in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by shelduck other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A050 Wigeon *Anas penelope***

**To maintain the favourable conservation condition of Wigeon in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by wigeon other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*



**A052 Teal *Anas crecca***

**To maintain the favourable conservation condition of Teal in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by teal other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A054 Pintail *Anas acuta***

**To maintain the favourable conservation condition of Pintail in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by pintail other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A056 Shoveler *Anas clypeata***

**To maintain the favourable conservation condition of Shoveler in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by shoveler other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A062 Scaup *Aythya marila***

**To maintain the favourable conservation condition of Scaup in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by scaup other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A137 Ringed Plover *Charadrius hiaticula***

To maintain the favourable conservation condition of Ringed Plover in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by ringed plover other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

For inspection purposes only.  
Consent of copyright owner required for any other use.

**A140 Golden Plover *Pluvialis apricaria***

To maintain the favourable conservation condition of Golden Plover in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by golden plover other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

For inspection purposes only.  
Consent of copyright owner required for any other use.

**A141 Grey Plover *Pluvialis squatarola***

**To maintain the favourable conservation condition of Grey Plover in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by grey plover other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A142 Lapwing *Vanellus vanellus***

To maintain the favourable conservation condition of Lapwing in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by lapwing other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

For inspection purposes only.  
Consent of copyright owner required for any other use.



**A143 Knot *Calidris canutus***

**To maintain the favourable conservation condition of Knot in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by knot other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A149 Dunlin *Calidris alpina***

**To maintain the favourable conservation condition of Dunlin in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by dunlin other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A156 Black-tailed Godwit *Limosa limosa***

**To maintain the favourable conservation condition of Black-tailed Godwit in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by black-tailed godwit other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A157 Bar-tailed Godwit *Limosa lapponica***

To maintain the favourable conservation condition of Bar-tailed Godwit in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by bar-tailed godwit other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

For inspection purposes only.  
Consent of copyright owner required for any other use.

**A160 Curlew *Numenius arquata***

**To maintain the favourable conservation condition of Curlew in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by curlew other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A162 Redshank *Tringa totanus***

**To maintain the favourable conservation condition of Redshank in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by redshank other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A164 Greenshank *Tringa nebularia***

**To maintain the favourable conservation condition of Greenshank in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by greenshank other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

**A179 Black-headed Gull *Chroicocephalus ridibundus***

To maintain the favourable conservation condition of Black-headed Gull in the River Shannon and River Fergus Estuaries SPA, which is defined by the following list of attributes and targets:

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Population trend	Percentage change	Long term population trend stable or increasing	Waterbird population trends are presented in part four of the conservation objectives supporting document
Distribution	Range, timing and intensity of use of areas	There should be no significant decrease in the range, timing or intensity of use of areas by black-headed gull other than that occurring from natural patterns of variation	As determined by regular low tide and other waterbird surveys. Waterbird distribution from the 2010/2011 waterbird survey programme is discussed in part five of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*



**A999 Wetlands**

**To maintain the favourable conservation condition of the wetland habitat in the River Shannon and River Fergus Estuaries SPA as a resource for the regularly-occurring migratory waterbirds that utilise it. This is defined by the following attribute and target:**

<b>Attribute</b>	<b>Measure</b>	<b>Target</b>	<b>Notes</b>
Wetland habitat area	hectares	The permanent area occupied by the wetland habitat should be stable and not significantly less than the area of 32,261ha, other than that occurring from natural patterns of variation	The wetland habitat area was estimated as 32,261ha using OSi data and relevant orthophotographs. For further information see part three of the conservation objectives supporting document

*For inspection purposes only.  
Consent of copyright owner required for any other use.*




**Legend**

SPA 004077

An Roinn Ealaíon, Oidhreacht agus Gaeltachta  
 Department of Arts, Heritage and the Gaeltacht

MAP 1:  
 RIVER SHANNON AND RIVER FERGUS ESTUARIES  
 CONSERVATION OBJECTIVES  
 SPA DESIGNATION


Map to be read in conjunction with the NPWS  
 Conservation Objectives Document.



Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).

Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaithe. Macasamhail d'ábhar na Suirbhéarachta Ordonáis le chead ón Rialtas (Ceadúnas Uimh. EN 0059208)

Site Code  
 SPA 004077  
 Version: CL 1.03 / LI 1.03  
 / KE 1.02  
 Map Version 1  
 Date: May 2012



002165

004077

For inspection purposes only.  
Consent of copyright owner required for any other use.

### Legend

- SPA 004077 River Shannon and River Fergus Estuaries
- SAC 002165 Lower River Shannon
- OSi Discovery Series County Boundaries



## **B6 Permits and Licences**

In 2000, the landowner Cussen & Co (Crane Hire) Limited, now CLIIB, was granted the original Waste Licence (W0082-01). In 2001 IPODEC Ireland Ltd. which was renamed Onyx and subsequently Veolia Environmental Services Ireland Ltd (VESI), acquired the Cussen waste business.

In November 2003 a revised Licence (W0082-02) was granted to VESI. In 2010, Greenstar Environmental Services (GES) acquired the trade and assets of VESI, which included the Ballykeefe facility and the name of the licensee was changed to GES. Starrus Eco Holdings Ltd acquired the trade and assets of GES and the Licence was transferred in March 2014.

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

## **B7 Connection to Foul Sewer**

It is a condition of the current planning permission 13/300 (Attachment B6) that a connection is made to the Limerick Main Drainage Scheme and is operated to the satisfaction of the planning authority in accordance with permissions 06/1394 and 08/2320. The landowner has engaged with the Water Services Authority in relation to the installation of a new foul sewer and connection to the Limerick Main Drainage Scheme and a copy of correspondence confirming this is enclosed in this Attachment.

*For inspection purposes only.  
Consent of copyright owner required for any other use.*

# CLLIB HOLDINGS LTD.

DOCK ROAD LIMERICK.

Greenstar Ltd,  
Dock Road,  
Limerick.

23/01/14


Dear Emma,

I refer to our conversation with regard to the sewer connection from the Greenstar facility on the Dock Road, Limerick to the waste water treatment plant.

At this time a wayleave has been granted across adjacent lands at the rear of your facility by Limerick City Council who are the owners of this land.

A survey of the ground and the route for the pipe was carried out last week and we are awaiting the results of this so we can proceed with putting this project out to tender asap.

Yours Sincerely,



William Cussen,

CLLIB Holdings Ltd.



Limerick County Council,  
Planning,  
County Hall,  
Dorradoyle,  
County Limerick.

2<sup>nd</sup> April 2014

RE: Application for a Licence – Greenstar Environmental Services Limited (In Receivership)  
Notice under section 87(1)(a) of the Act of 1992

Dear Sir / Madam,

Starrus Eco Holding Ltd, Dock Road, Limerick, is applying to the Environmental Protection Agency (EPA) for a Licence for its existing Materials Recovery Facility at Ballykeffe, Dock Road, Limerick. It is intended to increase the amount of waste accepted at the facility to 130,000 tonnes. Planning permission has been granted for the development (Ref 13/300). The classes and nature of the industrial emissions directive activities in accordance with the First Schedule to the Act of 1992 are:

**11.1 The recovery or disposal of waste in a facility, within the meaning of the Act of 1996, which facility is connected or associated with another activity specified in this Schedule in respect of which a licence or revised licence under Part IV is in force or in respect of which a licence under the said Part is or will be required.**

**11.4.(b) Recovery, or a mix of recovery and disposal, of non-hazardous waste with a capacity exceeding 75 tonnes per day involving one or more of the following activities, (other than activities to which the Urban Waste Water Treatment Regulations 2001 (S.I. No. 254 of 2001) apply):**

**(ii) pre-treatment of waste for incineration or co-incineration;**

An Environmental Impact Statement will be submitted to the Agency with the application. The Environmental Impact Statement, and any further information relating to the effects on the environment of the emissions from the activity, which may be furnished to the Agency in the course of the Agency's consideration of the application, will be available at the headquarters of the Agency.


A copy of the Environmental Impact Statement was submitted to Limerick County Council with the planning application.

A copy of the application for the licence may be inspected on the Agency's website or inspected at or obtained from the headquarters of the Agency as soon as is practicable after the receipt by the Agency of the application for the licence.

Yours sincerely

  
Jim O'Callaghan

120482201/JOC/KC c.c.  
c.c. Mr. Malcolm Dowling, Greenstar




**O' Callaghan Moran & Associates.**  
 Granary House, Rutland Street,  
 Cork, Ireland.  
 Tel. (021) 4321521 Fax. (021) 4321522  
 email : info@ocallaghanmoran.com

CLIENT  
**Greenstar Environmental Services Ltd**

details

Figure No.  
 001

This drawing is the property of O'Callaghan Moran & Associates and shall not be used, reproduced or disclosed to anyone without the prior written permission of O'Callaghan Moran & Associates and shall be returned upon request.

TITLE  
**SITE LOCATION**

SCALE	REV.
1:2500 A4	A



## **APPLICATION TO THE ENVIRONMENTAL PROTECTION AGENCY FOR A LICENCE**

Starrus Eco Holding Limited, Ballykeefe Townland, Dock Road, Limerick, is applying to the Environmental Protection Agency (EPA) for a Licence for its existing Materials Recovery Facility at Ballykeeffe Townland, Dock Road, Limerick. It is intended to increase the annual amount of waste accepted at the facility to 130,000 tonnes. The classes and nature of the industrial emissions directive activities in accordance with the First Schedule to the Act of 1992 are:

**11.1 The recovery or disposal of waste in a facility, within the meaning of the Act of 1996, which facility is connected or associated with another activity specified in this Schedule in respect of which a licence or revised licence under Part IV is in force or in respect of which a licence under the said Part is or will be required.**

**11.4.(b) Recovery, or a mix of recovery and disposal, of non-hazardous waste with a capacity exceeding 75 tonnes per day involving one or more of the following activities, (other than activities to which the Urban Waste Water Treatment Regulations 2001 (S.I. No. 254 of 2001) apply):**

**(ii) pre-treatment of waste for incineration or co-incineration;**

An Environmental Impact Statement will be submitted to the Agency with the application. A copy of this application, including the Environmental Impact Statement, as well such further information relating to the effects on the environment of the emissions from the activity which may be furnished to the Agency in the course of the Agency's consideration of the application, will be available at the headquarters of the Agency.

A copy of the Environmental Impact Statement has been submitted to Limerick County Council.

A copy of the application for the licence may be inspected on the Agency's website or inspected at or obtained from the headquarters of the Agency as soon as is practicable after the receipt by the Agency of the application for the licence.

Dated: 2<sup>nd</sup> April 2014

**COFFEE MORNING:** A coffee morning will take place in the Community Resource Centre in Ballybrown in aid of Millford Hospice on Friday, March 21 from 10.30 to 12.30 your support will be much appreciated the Resource Centre will be open from 9.30 to take in cakes etc.

**ASKEATON AFC:** Askeaton AFC committee meeting will be held on Monday

**TOURIST OFFICE:** The community information point at the tourist office is

School in UL and the business student Teachers' Association of Ireland. The Ms McMahon at a ceremony held in the University of Limerick on March 5.

advertising@limerickleader.ie | www.limerickleader.ie/ads | Tel: 087 274631/445/677 | Fax: 081 407422

# Classifieds

**Post to:**  
Limerick Leader  
107 Canal Street  
Limerick  
Co. Limerick

**Deadline:**  
Classifieds & Display Advertisements  
Monday 2:00 pm

**Property Management & Commercial Services:**  
Monday 2:00 pm

**State of Property/Job Vacancies:**  
Monday 2:00 pm

A complete range of services available to assist you in the sale or purchase of your property. We offer a free valuation service and a free estimate of the cost of repairs to your property.

**PROPERTY WA**

David  
**O' Driscoll**  
& A

For ready to go buyers in City Centre up to €150,000

Please call David O'Driscoll on 087 274631  
email: info@dodriscoll.com

**ECO ROOF**

Specialists in All Types of New and Old, We do It

Felting, Lathing, Tying, Slatting, Felt and Cladding

For a Great Quote Contact James 087 9688224; Robert 087 9688225  
Office 061 518318

**Small Ad:**  
Private training and coaching available for all levels of fitness and health. Contact: 087 274631

**ACCOMMODATION**

**APARTMENT TO LET**  
Wolfe Tone St - one bed self contained apartment, fully renovated, excellent condition throughout, very private, all mod cons. Call 087 2618863

**CITY CENTRE 3 Bed house, gfch, fully furnished, all mod cons, parking. 600pm Call 086 8342829**

**PARKLAND APTS**  
Dublin Rd, two apartments all mod cons. Private parking. Contact 087 2747010

**ACCOUNTING & FINANCIAL SERVICES**  
ACCOUNTANT

**DOGS**

**Dog training** Residential guaranteed dog training in Obedience and behaviour problems for all breeds, ie pulling on lead, house-training, aggression, recall, chewing, jumping up etc. Assistance/service dogs and training available for Autism, Diabetes, Hearing, Seizures and Mobility. Over 20 years experience, having trained in the UK, USA and Europe. Qualified master dog

**FARM & GARDEN**

**BEECH, LAURAL**  
Griselinia, Whitethorn, Blackthorn, Hazel, Dogwood, Hornbeam and all other Hedging plants. Also 6 - 8ft Trees only 15 - 20. Special Offer, Best Prices. Delivery arranged. Kearney's Nursery, Galbally 086 862 6736  
www.irelandtrees.com

**CONWAY PLANT HIRE** Land Reclamation, Dykes, Drains, Ditch Stretching and Cutting. All concrete work and re seeding. Excavators, Dozers,

**INDUST & COMMERCIAL PROP**

**FOR SALE** Corner Retail Unit. 530sqft. The Milk Market, Carr Street, Limerick. Contact Rooney Auctioneers 061 413511

**LIVESTOCK WANTED**

**WANTED HEIFERS**  
Bullocks & Young Bulls, fit for slaughter. Contact Conor Kennedy Fedamore, at John Kelly Meats Ltd. 087-4155191.

**PERSONAL**

**IRELAND**

**ADKI**

Property restoration including emergency response, fire/smoke damage, flood/water damage and property repair

Call: 1890-354-354


**LISTON Auctioneers**

**twitter**  
@limerick\_leader

**PUBLIC NOTICES**

An Chuir Chuarda The Circuit Ct  
EPA Export 16-04-2014:23:36:08

# Wool Centre



**9.00**

Program  
uttnohole  
Beginners

**€155.00**

14 Stitch Program  
4 Step Buttonhole  
Portable for Sewing Classes

## FREE SEWING KIT

## MADE INS WELCOME MONSTRATIONS DAILY

SERVICE ON ALL MAKES OF MACHINES

14, OFF WILLIAM STREET, LIMERICK  
061 414829

## Wm Fitzgerald Skip Hire Ltd

[www.fitzgeraldskiphire.com](http://www.fitzgeraldskiphire.com)

& start your Spring Clean today  
DIRTY LOCAL - CALL 061 414200



12 YD BULKER

Local Family Run

Reliable



8 YD STANDARD

FAMILY owned and run

20 years

1 fastest skip service



6 YD MEDIUM

Ennis, Rathbane North, Limerick.

061-414200 & 061-414200

Email: [info@fitzgeraldskiphire.com](mailto:info@fitzgeraldskiphire.com)

## gham Holiday Homes

Villerby & BK agents

Large selection of New & Used mobile  
homes in stock at Ennis Road, Limerick.  
Monday - Saturday 061 451009

2 bedroom  
homes  
on site  
see and  
buy point  
€10,000.



Viewing by appointment.

Contact Brian 061 451009

# WANTED

Friesian Bulls -  
table for feeding

## APPLICATION TO THE ENVIRONMENTAL PROTECTION AGENCY FOR A LICENCE

Starrus Eco Holdings Limited, Ballykeefe Townland, Dock Road, Limerick, is applying to the Environmental Protection Agency (EPA) for a Licence for its existing Materials Recovery Facility at Ballykeefe Townland, Dock Road, Limerick. It is intended to increase the annual amount of waste acceptable at the facility to 130,000 tonnes. The classes and nature of the industrial emissions; directive activities in accordance with the First Schedule to the Act of 1992 are:

11.1 The recovery or disposal of waste in a facility, within the meaning of the Act of 1996, which facility is connected or associated with another activity specified in this Schedule in respect of which a licence or revised licence under Part IV is in force or in respect of which a licence under the said Part is or will be required.

11.4.(b) Recovery, or a mix of recovery and disposal, of non-hazardous waste with a capacity exceeding 75 tonnes per day involving one or more of the following activities, (other than activities to which the Urban Waste Water Treatment Regulations 2001 (S.I. No. 254 of 2001) apply):

(ii) pre-treatment of waste for incineration or co-incineration;

An Environmental Impact Statement will be submitted to the Agency with the application. A copy of this application, including the Environmental Impact Statement, as well as such further information relating to the effects on the environment of the emissions from the activity which may be furnished to the Agency in the course of the Agency's consideration of the application, will be available at the headquarters of the Agency.

A copy of the Environmental Impact Statement has been submitted to Limerick County Council.

A copy of the application for the licence may be inspected on the Agency's website or inspected at or obtained from the headquarters of the Agency as soon as is practicable after the receipt by the Agency of the application for the licence.

This notice refers to the same remediation project as previously advertised (W0281-01), which is required to be re-advertised due to changes in the legislation. The Waste Licence Application is required to be regarded to an Industrial Emissions Licence Application

## APPLICATION TO THE ENVIRONMENTAL PROTECTION AGENCY FOR A LICENCE

NOTICE is hereby given in accordance with Articles 5 and 6 of the Environmental Protection Agency (Industrial Emissions) (licensing) Regulations, 2013 (as amended) (SI No. 137 of 2013) that "Bord Gáis Éireann" having its principal offices at Gasworks Road, Cork, will apply for an Industrial Emissions Licence to the Environmental Protection Agency by the prescribed date in respect of land bounded by Dock Road to the north west, O'Curry Street to the north east and housing and light industry to the south east and south west in Limerick, otherwise known as "Former Limerick Gasworks" - National Grid Reference:-R 5694E, 5656N The works will comprise the remediation (clean up) of the Former Limerick Gasworks, involving pumping and treating of groundwater and stabilising soils, with associated recovery and disposal of non-hazardous and hazardous contaminated soils.

An Environmental Impact Statement has been previously submitted to the Agency in relation to a Waste Licence application (W0281-01).

The classes of activity as per the First Schedule of the EPA Acts 1992 to 2013 are:

11.1- The recovery or disposal of waste in a facility, within the meaning of the Act of 1996, which facility is connected or associated with another activity specified in this schedule in respect of which a licence or revised licence under Part IV is in force or in respect of which a licence under the said Part is or will be required.

11.2- Disposal or recovery of hazardous waste with a capacity exceeding 10 tonnes per day involving one or more of the following activities:

(b) physio-chemical treatment;

A copy of the licence application (W0281-01), Environmental Impact Statement and any such further information relating to the effects on the environment of the emissions from the activity which may be furnished to the Agency in the course of the Agency's consideration of the application will be available at the headquarters of the Agency. A copy of the Environmental Impact Statement has been submitted to the Limerick City Council planning authority.

A copy of the licence application may be inspected on the Agency's website or inspected at or obtained from the headquarters of the Agency as soon as is practicable after the receipt by the Agency of the application for the licence.

## APPLICATION TO THE ENVIRONMENTAL PROTECTION AGENCY FOR A LICENCE

Starrus Eco Holdings Limited, Ballykeefe Townland, Dock Road, Limerick, is applying to the Environmental Protection Agency (EPA) for a Licence for its existing Materials Recovery Facility at Ballykeefe Townland, Dock Road, Limerick. It is intended to increase the annual amount of waste acceptable at the facility to 130,000 tonnes. The classes and nature of the industrial emissions directive activities in accordance with the First Schedule to the Act of 1992 are:

**11.1 The recovery or disposal of waste in a facility, within the meaning of the Act of 1996, which facility is connected or associated with another activity specified in this Schedule in respect of which a licence or revised licence under Part IV is in force or in respect of which a licence under the said Part is or will be required.**

**11.4.(b) Recovery, or a mix of recovery and disposal, of non-hazardous waste with a capacity exceeding 75 tonnes per day involving one or more of the following activities, (other than activities to which the Urban Waste Water Treatment Regulations 2001 (S.I. No. 254 of 2001) apply):**

**(ii) pre-treatment of waste for incineration or co-incineration;**

An Environmental Impact Statement will be submitted to the Agency with the application. A copy of this application, including the Environmental Impact Statement, as well such further information relating to the effects on the environment of the emissions from the activity which may be furnished to the Agency in the course of the Agency's consideration of the application, will be available at the headquarters of the Agency.

A copy of the Environmental Impact Statement has been submitted to Limerick County Council.

A copy of the application for the licence may be inspected on the Agency's website or inspected at or obtained from the headquarters of the Agency as soon as is practicable after the receipt by the Agency of the application for the licence.