

APPENDIX A
Ballyman Glen SAC Site Synopsis Report

For inspection purposes only.
Consent of copyright owner required for any other use.

For inspection purposes only.
Consent of copyright owner required for any other use.



Site Name: Ballyman Glen SAC

Site Code: 000713

Ballyman Glen is situated approximately 3 km north of Enniskerry and straddles the County boundary between Dublin and Wicklow. It is orientated in an east-west direction with a stream running through the centre. The glen is bounded mostly by steeply sloping pasture with Gorse (*Ulex europaeus*) and areas of wood and scrub.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes):

[7220] Petrifying Springs*

[7230] Alkaline Fens

Ballyman Glen contains a small strip of alkaline fen which is associated with petrifying spring/seepage areas that have given rise to thick deposits of marl. The vegetation of the main part of the fen is dominated by Greater Tussock-sedge (*Carex paniculata*), Tall Fescue (*Festuca arundinacea*), butterworts (*Pinguicula vulgaris* and *P. lusitanica*), Black Bog-rush (*Schoenus nigricans*) and Broad-leaved Cottongrass (*Eriophorum latifolium*). The site is particularly notable for its orchids, with species including Early Marsh-orchid (*Dactylorhiza incarnata*), Narrow-leaved Marsh-orchid (*D. traunsteineri*) and Marsh Helleborine (*Epipactis palustris*) occurring. In addition, twenty species of sedge have been recorded in the area, including the scarce Long-stalked Yellow-sedge (*Carex lepidocarpa*). The fen area is being invaded by Downy Birch (*Betula pubescens*).

Associated with the fen, and also with the woodland elsewhere in the site, are petrifying springs. These lime-encrusted seepage areas are rich in bryophytes including such diagnostic species as *Cratoneuron commutatum* and *C. filicinum*.

Wet woodland and scrub occur along the margins of the stream for most of the length of the glen, extending outwards in areas to create inaccessible and species-rich patches of woodland. The canopy is dominated by Alder (*Alnus glutinosa*), willow (*Salix* spp.) and Ash (*Fraxinus excelsior*). The woodland has a dense shrub layer which includes Hawthorn (*Crataegus monogyna*) and Spindle (*Euonymus europaeus*), and a diverse ground flora with Marsh Hawk's-beard (*Crepis paludosa*), Sanicle (*Sanicula europaea*), Herb-Robert (*Geranium robertianum*), Bugle (*Ajuga reptans*), horsetails (*Equisetum* spp.), Meadowsweet (*Filipendula ulmaria*) and some sedges (*Carex* spp.). Areas of marsh are found in the wetter areas by the stream, particularly at the western end of the site.

There is an area of broadleaved woodland on the steeper southern slopes of the glen. Common species occurring here are Ash and Sycamore (*Acer pseudoplatanus*), with Bramble (*Rubus fruticosus* agg.) colonizing the more open areas.

An area of land that slopes towards the fen has been used as a landfill site for domestic refuse. The site is also used for a clay pigeon shoot and shattered clay pigeons are scattered throughout the area.

The fen vegetation at this site is well developed, with an unusually large number of sedge species present. The presence of alkaline fen and of petrifying spring/seepage areas is also particularly notable, as these habitats are listed, the latter with priority status, on Annex I of the E.U. Habitats Directive. Fens are rare in Wicklow and Dublin, and this is one of only two sites in Wicklow for the Narrow-leaved Marsh-orchid.

For inspection purposes only.
Consent of copyright owner required for any other use.