




CONSULTANTS IN ENGINEERING,
ENVIRONMENTAL SCIENCE & PLANNING

APPENDIX 4

European Site Synopses

*For inspection purposes only.
Consent of copyright owner required for any other use.*





Site Name: Lower River Shannon SAC

Site Code: 002165

This very large site stretches along the Shannon valley from Killaloe in Co. Clare to Loop Head/ Kerry Head, a distance of some 120 km. The site thus encompasses the Shannon, Feale, Mulkear and Fergus estuaries, the freshwater lower reaches of the River Shannon (between Killaloe and Limerick), the freshwater stretches of much of the Feale and Mulkear catchments and the marine area between Loop Head and Kerry Head. Rivers within the sub-catchment of the Feale include the Galey, Smearlagh, Oolagh, Allaughaun, Owveg, Clydagh, Caher, Breanagh and Glenacarne. Rivers within the sub-catchment of the Mulkear include the Killeenagarriff, Annagh, Newport, the Dead River, the Bilboa, Glashacloonaraveela, Gortnageragh and Cahernahallia.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes):

- [1110] Sandbanks
- [1130] Estuaries
- [1140] Tidal Mudflats and Sandflats
- [1150] Coastal Lagoons*
- [1160] Large Shallow Inlets and Bays
- [1170] Reefs
- [1220] Perennial Vegetation of Stony Banks
- [1230] Vegetated Sea Cliffs
- [1310] *Salicornia* Mud
- [1330] Atlantic Salt Meadows
- [1410] Mediterranean Salt Meadows
- [3260] Floating River Vegetation
- [6410] *Molinia* Meadows
- [91E0] Alluvial Forests*
- [1029] Freshwater Pearl Mussel (*Margaritifera margaritifera*)
- [1095] Sea Lamprey (*Petromyzon marinus*)
- [1096] Brook Lamprey (*Lampetra planeri*)
- [1099] River Lamprey (*Lampetra fluviatilis*)
- [1106] Atlantic Salmon (*Salmo salar*)
- [1349] Bottle-nosed Dolphin (*Tursiops truncatus*)
- [1355] Otter (*Lutra lutra*)

The Shannon and Fergus Rivers flow through Carboniferous limestone as far as Foynes, but west of Foynes Namurian shales and flagstones predominate (except at Kerry Head, which is formed from Old Red Sandstone). The eastern sections of the Feale catchment flow through Namurian rocks and the western stretches through Carboniferous limestone. The Mulkear flows through Lower Palaeozoic rocks in the upper reaches before passing through Namurian rocks, followed by Lower Carboniferous shales and Carboniferous limestone. The Mulkear River itself, immediately north of Pallas Green, passes through an area of Rhyolites, Tuffs and Agglomerates.

The Shannon and Fergus Estuaries form the largest estuarine complex in Ireland. They form a unit stretching from the upper tidal limits of the Shannon and Fergus Rivers to the mouth of the Shannon Estuary (considered to be a line across the narrow strait between Kilcredaun Point and Kilconly Point). Within this main unit there are several tributaries with their own 'sub-estuaries' e.g. the Deel River, Mulkear River, and Maigue River. To the west of Foynes, a number of small estuaries form indentations in the predominantly hard coastline, namely Poulnasherry Bay, Ballylongford Bay, Clonderalaw Bay and the Feale or Cashen River estuary.

Both the Fergus and inner Shannon Estuaries feature vast expanses of intertidal mudflats, often fringed with saltmarsh vegetation. The smaller estuaries also feature mudflats, but have their own unique characteristics, e.g. Poulnasherry Bay is stony and unusually rich in species and biotopes. Plant species are typically scarce on the mudflats, although there are some eelgrass (*Zostera* spp.) beds and patches of green algae (e.g. *Ulva* sp. and *Enteromorpha* sp.). The main macro-invertebrate community which has been noted from the inner Shannon and Fergus estuaries is a *Macoma-Scrobicularia-Nereis* community.

In the transition zone between mudflats and saltmarsh, specialised colonisers of mud predominate. For example, swards of Common Cord-grass (*Spartina anglica*) frequently occur in the upper parts of the estuaries. Less common are swards of Glasswort (*Salicornia europaea* agg.). In the innermost parts of the estuaries, the tidal channels or creeks are fringed with species such as Common Reed (*Phragmites australis*) and club-rushes (*Scirpus maritimus*, *S. tabernaemontani* and *S. triquetrus*). In addition to the nationally rare Triangular Club-rush (*Scirpus triqueter*), two scarce species are found in some of these creeks (e.g. Ballinacurra Creek): Lesser Bulrush (*Typha angustifolia*) and Summer Snowflake (*Leucojum aestivum*).

Saltmarsh vegetation frequently fringes the mudflats. Over twenty areas of estuarine saltmarsh have been identified within the site, the most important of which are around the Fergus estuary and at Ringmoylan Quay. The dominant type of saltmarsh present is Atlantic salt meadow occurring over mud. Characteristic species occurring include Common Saltmarsh-grass (*Puccinellia maritima*), Sea Aster (*Aster tripolium*), Thrift (*Armeria maritima*), Sea-milkwort (*Glaux maritima*), Sea Plantain (*Plantago maritima*), Red Fescue (*Festuca rubra*), Creeping Bent (*Agrostis stolonifera*), Saltmarsh Rush (*Juncus gerardi*), Long-bracted Sedge (*Carex extensa*), Lesser Sea-spurrey

(*Spergularia marina*) and Sea Arrowgrass (*Triglochin maritima*). Areas of Mediterranean salt meadows, characterised by clumps of Sea Rush (*Juncus maritimus*) occur occasionally. Two scarce species are found on saltmarshes in the vicinity of the Fergus estuary: a type of robust saltmarsh-grass (*Puccinellia foucaudii*), sometimes placed within the species Common Saltmarsh-grass (*P. maritima*) and Hard-grass (*Parapholis strigosa*).

Saltmarsh vegetation also occurs around a number of lagoons within the site, two of which have been surveyed as part of a National Inventory of Lagoons. Cloonconeen Pool (4-5 ha) is a natural sedimentary lagoon impounded by a low cobble barrier. Seawater enters by percolation through the barrier and by overwash. This lagoon represents a type which may be unique to Ireland since the substrate is composed almost entirely of peat. The adjacent shore features one of the best examples of a drowned forest in Ireland. Aquatic vegetation in the lagoon includes typical species such as Beaked Tasselweed (*Ruppia maritima*) and green algae (*Cladophora* sp.). The fauna is not diverse, but is typical of a high salinity lagoon and includes six lagoon specialists (*Hydrobia ventrosa*, *Cerastoderma glaucum*, *Lekanesphaera hookeri*, *Palaemonetes varians*, *Sigara stagnalis* and *Enochrus bicolor*). In contrast, Shannon Airport Lagoon (2 ha) is an artificial saline lake with an artificial barrier and sluiced outlet. However, it supports two Red Data Book species of stonewort (*Chara canescens* and *Chara cf. connivens*).

Most of the site west of Kilcredaun Point/Kilconly Point is bounded by high rocky sea cliffs. The cliffs in the outer part of the site are sparsely vegetated with lichens, Red Fescue, Sea Beet (*Beta vulgaris* subsp. *maritima*), Sea Campion (*Silene vulgaris* subsp. *maritima*), Thrift and plantains (*Plantago* spp.). A rare endemic type of sea-lavender, *Limonium recurvum* subsp. *pseudotranswallianum*, occurs on cliffs near Loop Head. Cliff-top vegetation usually consists of either grassland or maritime heath. The boulder clay cliffs further up the estuary tend to be more densely vegetated, with swards of Red Fescue and species such as Kidney Vetch (*Anthyllis vulneraria*) and Common Bird's-foot-trefoil (*Lotus corniculatus*).

The site supports an excellent example of a large shallow inlet and bay. Littoral sediment communities in the mouth of the Shannon Estuary occur in areas that are exposed to wave action and also in areas extremely sheltered from wave action. Characteristically, exposed sediment communities are composed of coarse sand and have a sparse fauna. Species richness increases as conditions become more sheltered. All shores in the site have a zone of sand hoppers at the top, and below this each of the shores has different characteristic species giving a range of different shore types.

The intertidal reefs in the Shannon Estuary are exposed or moderately exposed to wave action and subject to moderate tidal streams. Known sites are steeply sloping and show a good zonation down the shore. Well developed lichen zones and littoral reef communities offering a high species richness in the sublittoral fringe and strong populations of the Purple Sea Urchin *Paracentrotus lividus* are found. The communities found are tolerant to sand scour and tidal streams. The infralittoral reefs range from sloping platforms with some vertical steps, to ridged bedrock with

gullies of sand between the ridges, to ridged bedrock with boulders or a mixture of cobbles, gravel and sand. Kelp is very common to about 18 m. Below this it becomes rare and the community is characterised by coralline crusts and red foliose algae.

Other coastal habitats that occur within the site include stony beaches and bedrock shores (these support a typical zonation of seaweeds such as *Fucus* spp., *Ascophyllum nodosum* and kelps), shingle beaches (with species such as Sea Beet, Sea Mayweed - *Matricaria maritima*, Sea Campion and Curled Dock - *Rumex crispus*), sandbanks which are slightly covered by sea water at all times (e.g. in the area from Kerry Head to Beal Head) and sand dunes (a small area occurs at Beal Point, where Marram - *Ammophila arenaria* is the dominant species).

Freshwater rivers have been included in the site, most notably the Feale and Mulkear catchments, the Shannon from Killaloe to Limerick (along with some of its tributaries, including a short stretch of the Kilmastulla River), the Fergus up as far as Ennis, and the Cloon River. These systems are very different in character: the Shannon is broad, generally slow flowing and naturally eutrophic; the Fergus is smaller and alkaline; while the narrow, fast flowing Cloon is acid in nature. The Feale and Mulkear catchments exhibit all the aspects of a river from source to mouth. Semi-natural habitats, such as wet grassland, wet woodland and marsh occur by the rivers, but improved grassland is the most common habitat type. One grassland type of particular conservation significance, *Molinia* meadows, occurs in several parts of the site and the examples at Worldsend on the River Shannon are especially noteworthy. Here are found areas of wet meadow dominated by rushes (*Juncus* spp.) and sedges (*Carex* spp.), and supporting a diverse and species-rich vegetation, including such uncommon species as Blue-eyed Grass (*Sisyrinchium bermudiana*) and Pale Sedge (*C. pallescens*).

Floating river vegetation characterised by species of water-crowfoot (*Ranunculus* spp.), pondweeds (*Potamogeton* spp.) and the moss *Fontinalis antipyretica* are present throughout the major river systems within the site. The rivers contain an interesting bryoflora with *Schistidium alpicola* var. *alpicola* recorded from in-stream boulders on the Bilboa, new to Co. Limerick.

Alluvial woodland occurs on the banks of the Shannon and on islands in the vicinity of the University of Limerick. The woodland is up to 50 m wide on the banks and somewhat wider on the largest island. The most prominent woodland type is gallery woodland where White Willow (*Salix alba*) dominates the tree layer with occasional Alder (*Alnus glutinosa*). The shrub layer consists of various willow species with Rusty Willow (*Salix cinerea* ssp. *oleifolia*) and what appear to be hybrids of *S. alba* x *S. viminalis*. The herbaceous layer consists of tall perennial herbs. A fringe of bulrush (*Typha* sp.) occurs on the river side of the woodland. On slightly higher ground above the wet woodland and on the raised embankment remnants of mixed oak-ash-alder woodland occur. These are poorly developed and contain numerous exotic species but locally there are signs that it is invading open grassland. Alder is the principal tree species, with occasional Pedunculate Oak (*Quercus robur*), elm (*Ulmus glabra* and *U. procera*), Hazel (*Corylus avellana*), Hawthorn (*Crataegus monogyna*) and

the shrubs Guelder-rose (*Viburnum opulus*) and willows. The ground flora is species-rich.

While woodland is infrequent within the site, however Cahiracon Wood contains a strip of old oak woodland. Sessile Oak (*Q. petraea*) forms the canopy, with an understorey of Hazel and Holly (*Ilex aquifolium*). Great Wood-rush (*Luzula sylvatica*) dominates the ground flora. Less common species present include Great Horsetail (*Equisetum telmateia*) and Pendulous Sedge (*Carex pendula*).

In the low hills to the south of the Slievefelim Mountains, the Cahernahallia River cuts a valley through the Upper Silurian rocks. For approximately 2 km south of Cappagh Bridge at Knockanavar, the valley sides are wooded. The woodland consists of birch (*Betula* spp.), Hazel, oak, Rowan (*Sorbus aucuparia*), some Ash (*Fraxinus excelsior*) and willow (*Salix* spp.). Most of the valley is not grazed by stock, and as a result the trees are regenerating well. The ground flora features prominent Great wood-rush and Bilberry (*Vaccinium myrtillus*), along with a typical range of woodland herbs. Bracken (*Pteridium aquilinum*) is a feature in areas where there is more light available.

The valley sides of the Bilboa and Gortnageragh Rivers on higher ground north-east of Cappamore, support patches of semi-natural broadleaf woodland dominated by Ash, Hazel, oak and birch. There is a good scrub layer with Hawthorn, willow, Holly and Blackthorn (*Prunus spinosa*) common. The herb layer in these woodlands is often open, with a typically rich mixture of woodland herbs and ferns. Moss species diversity is high. The woodlands are ungrazed. The Hazel is actively coppiced in places.

There is a small area of actively regenerating cut-away raised bog at Ballyrorheen. It is situated approximately 5 km north-west of Cappamore in Co. Limerick. The bog contains some wet areas with good cover of bog mosses (*Sphagnum* spp.). Species of particular interest include Cranberry (*Vaccinium oxycoccos*) and White Sedge (*Carex curta*), along with two regionally rare mosses, including the bog moss *S. fimbriatum*. The site is being invaded by Downy Birch (*Betula pubescens*) scrub woodland. Both commercial forestry and the spread of Rhododendron (*Rhododendron ponticum*) has greatly reduced the overall value of the site.

A number of plant species that are listed in the Irish Red Data Book occur within the site, and several of these are protected under the Flora (Protection) Order, 1999. These include Triangular Club-rush (*Scirpus triquetrus*), a species which is only found in Ireland only in the Shannon Estuary, where it borders creeks in the inner estuary. Opposite-leaved Pondweed (*Groenlandia densa*) is found in the Shannon where it passes through Limerick City, while Meadow Barley (*Hordeum secalinum*) is abundant in saltmarshes at Ringmoylan and Mantlehill. Hairy Violet (*Viola hirta*) occurs in the Askeaton/Foynes area. Golden Dock (*Rumex maritimus*) is noted as occurring in the River Fergus estuary. Finally, Bearded Stonewort (*Chara canescens*), a brackish water specialist, and Convergent Stonewort (*Chara connivens*) are both found in Shannon Airport Lagoon.

Overall, the Shannon and Fergus Estuaries support the largest numbers of wintering waterfowl in Ireland. The highest count in 1995-96 was 51,423 while in 1994-95 it was 62,701. Species listed on Annex I of the E.U. Birds Directive which contributed to these totals include: Great Northern Diver (3; 1994/95), Whooper Swan (201; 1995/96), Pale-bellied Brent Goose (246; 1995/96), Golden Plover (11,067; 1994/95) and Bar-tailed Godwit (476; 1995/96). In the past, three separate flocks of Greenland White-fronted Goose were regularly found, but none were seen in 1993/94.

Other wintering waders and wildfowl present include Greylag Goose (216; 1995/96), Shelduck (1,060; 1995/96), Wigeon (5,976; 1995/96), Teal (2,319; 1995-96), Mallard (528; 1995/96), Pintail (45; 1995/96), Shoveler (84; 1995/96), Tufted Duck (272; 1995/96), Scaup (121; 1995/96), Ringed Plover (240; 1995/96), Grey Plover (750; 1995/96), Lapwing (24,581; 1995/96), Knot (800; 1995/96), Dunlin (20,100; 1995/96), Snipe (719; 1995/96), Black-tailed Godwit (1,062; 1995/96), Curlew (1,504; 1995/96), Redshank (3,228; 1995/96), Greenshank (36; 1995/96) and Turnstone (107; 1995/96). A number of wintering gulls are also present, including Black-headed Gull (2,216; 1995/96), Common Gull (366; 1995/96) and Lesser Black-backed Gull (100; 1994/95). This is the most important coastal site in Ireland for a number of the waders including Lapwing, Dunlin, Snipe and Redshank. It also provides an important staging ground for species such as Black-tailed Godwit and Greenshank.

A number of species listed on Annex I of the E.U. Birds Directive breed within the site. These include Peregrine Falcon (2-3 pairs), Sandwich Tern (34 pairs on Rat Island, 1995), Common Tern (15 pairs: 2 on Sturamus Island and 13 on Rat Island, 1995), Chough (14-41 pairs, 1992) and Kingfisher. Other breeding birds of note include Kittiwake (690 pairs at Loop Head, 1987) and Guillemot (4,010 individuals at Loop Head, 1987).

There is a resident population of Bottle-nosed Dolphin in the Shannon Estuary. This is the only known resident population of this E.U. Habitats Directive Annex II species in Ireland. The population is estimated (in 2006) to be 140 ± 12 individuals. Otter, a species also listed on Annex II of this Directive, is commonly found on the site.

Five species of fish listed on Annex II of the E.U. Habitats Directive are found within the site. These are Sea Lamprey (*Petromyzon marinus*), Brook Lamprey (*Lampetra planeri*), River Lamprey (*Lampetra fluviatilis*), Twaite Shad (*Allosa fallax fallax*) and Salmon (*Salmo salar*). The three lampreys and Salmon have all been observed spawning in the lower Shannon or its tributaries. The Fergus is important in its lower reaches for spring salmon, while the Mulkear catchment excels as a grilse fishery, though spring fish are caught on the actual Mulkear River. The Feale is important for both types. Twaite Shad is not thought to spawn within the site. There are few other river systems in Ireland which contain all three species of lamprey.

Two additional fish species of note, listed in the Irish Red Data Book, also occur, namely Smelt (*Osmerus eperlanus*) and Pollan (*Coregonus autumnalis pollan*). Only the former has been observed spawning in the Shannon.

Freshwater Pearl Mussel (*Margaritifera margaritifera*), a species listed on Annex II of the E.U. Habitats Directive, occurs abundantly in parts of the Cloon River.

There is a wide range of land uses within the site. The most common use of the terrestrial parts is grazing by cattle, and some areas have been damaged through over-grazing and poaching. Much of the land adjacent to the rivers and estuaries has been improved or reclaimed and is protected by embankments (especially along the Fergus estuary). Further, reclamation continues to pose a threat, as do flood relief works (e.g. dredging of rivers). Gravel extraction poses a major threat on the Feale.

In the past, cord-grass (*Spartina* sp.) was planted to assist in land reclamation. This has spread widely, and may oust less vigorous colonisers of mud and may also reduce the area of mudflat available to feeding birds.

Domestic and industrial wastes are discharged into the Shannon, but water quality is generally satisfactory, except in the upper estuary where it reflects the sewage load from Limerick City. Analyses for trace metals suggest a relatively clean estuary with no influences of industrial discharges apparent. Further industrial development along the Shannon and water polluting operations are potential threats.

Fishing is a main tourist attraction on the Shannon and there are a large number of angler associations, some with a number of beats. Fishing stands and styles have been erected in places. The River Feale is a designated Salmonid Water under the E.U. Freshwater Fish Directive. Other uses of the site include commercial angling, oyster farming, boating (including dolphin-watching trips) and shooting. Some of these may pose threats to the birds and dolphins through disturbance. Specific threats to the dolphins include underwater acoustic disturbance, entanglement in fishing gear and collisions with fast moving craft.

This site is of great ecological interest as it contains a high number of habitats and species listed on Annexes I and II of the E.U. Habitats Directive, including the priority habitats lagoon and alluvial woodland, the only known resident population of Bottle-nosed Dolphin in Ireland and all three Irish lamprey species. A good number of Red Data Book species are also present, perhaps most notably the thriving populations of Triangular Club-rush. A number of species listed on Annex I of the E.U. Birds Directive are also present, either wintering or breeding. Indeed, the Shannon and Fergus Estuaries form the largest estuarine complex in Ireland and support more wintering wildfowl and waders than any other site in the country. Most of the estuarine part of the site has been designated a Special Protection Area (SPA), under the E.U. Birds Directive, primarily to protect the large numbers of migratory birds present in winter.

SITE SYNOPSIS

SITE NAME: KERRY HEAD SPA

SITE CODE: 004189

Kerry Head SPA is situated on the south side of the mouth of the River Shannon in north Co. Kerry. It encompasses the sea cliffs from just west of Ballyheigue, around the end of Kerry Head to the west and north-eastwards as far as Kilmore. The site includes the sea cliffs and land adjacent to the cliff edge. The high water mark forms the seaward boundary. Most of the site is underlain by Devonian siltstones, sandstones and mudstones; a small section of the site has rocks of Carboniferous age.

The site is a Special Protection Area (SPA) under the E.U. Birds Directive, of special conservation interest for the following species: Chough and Fulmar.

The site supports an internationally important population of breeding Chough, a Red Data Book species that is listed on Annex I of the E.U. Birds Directive; 32 breeding pairs were recorded from the site in the 1992 survey and 30 in the 2002/03 survey. In addition, a flock of 20 birds was noted on the northern coast of the site during the latter survey. The site is of particular note for the density of breeding pairs found.

The site also supports a nationally important population of Fulmar (421 pairs), as well as a small population of Shag (8 pairs) – all seabird data from 2000. The site is also used by Peregrine (2 pairs in 2002).

Kerry Head SPA is one of the most important sites in the country for Chough. It also supports a population of Fulmar of national importance. The presence of Chough and Peregrine, both species that are listed on Annex I of the E.U. Birds Directive, is of particular significance

20.1.2015

SITE SYNOPSIS

SITE NAME: STACK'S TO MULLAGHAREIRK MOUNTAINS, WEST LIMERICK HILLS AND MOUNT EAGLE SPA

SITE CODE: 004161

The Stack's to Mullaghareirk Mountains, West Limerick Hills and Mount Eagle SPA is a very large site centred on the borders between the counties of Cork, Kerry and Limerick. The site is skirted by the towns of Newcastle West, Ballydesmond, Castleisland, Tralee and Abbeyfeale. The mountain peaks included in the site are not notably high or indeed pronounced, the highest being at Knockfeha (451 m). Other mountains included are Mount Eagle, Knockanefune, Garraunbaun, Taur, Rock Hill, Knockacummer, Mullaghmuish, Knight's Mt, Ballincollig Hill, Beennageeha Mt, Sugar Hill, Knockanimpuba and Knockathea, amongst others. Many rivers rise within the site, notably the Blackwater, Owentaraglin, Owenkeal, Glenlara, Feale, Clydagh, Allaghaun, Allow, Oolagh, Galey and Smerlagh.

The site consists of a variety of upland habitats, though almost half is afforested. The coniferous forests include first and second rotation plantations, with both pre-thicket and post-thicket stands present. Substantial areas of clear-fell are also present at any one time. The principal tree species present are Sitka Spruce (*Picea sitchensis*) and Lodgepole Pine (*Pinus contorta*). A substantial part (28%) of the site is unplanted blanket bog and heath, with both wet and dry heath present. The vegetation of these habitats is characterised by such species as Ling Heather (*Calluna vulgaris*), Bilberry (*Vaccinium myrtillus*), Common Cottongrass (*Eriophorum angustifolium*), Hare's-tail Cottongrass (*Eriophorum vaginatum*), Deergrass (*Scirpus cespitosus*) and Purple Moor-grass (*Molinia caerulea*). The remainder of the site is mostly rough grassland that is used for hill farming. This varies in composition and includes some wet areas with rushes (*Juncus* spp.) and some areas subject to scrub encroachment.

The site is a Special Protection Area (SPA) under the E.U. Birds Directive, of special conservation interest for Hen Harrier.

This SPA is a stronghold for Hen Harrier and supports the largest concentration of the species in the country. A survey in 2005 recorded 45 pairs, which represents over 20% of the all-Ireland total. A similar number of pairs had been recorded in the 1998-2000 period. The mix of forestry and open areas provides optimum habitat conditions for this rare bird, which is listed on Annex I of the E.U. Birds Directive. The early stages of new and second-rotation conifer plantations are the most frequently used nesting sites, though some pairs may still nest in tall heather of unplanted bogs and heath. Hen Harriers will forage up to c. 5 km from the nest site, utilising open bog and moorland, young conifer plantations and hill farmland that is not too rank. Birds will often forage in openings and gaps within forests. In Ireland, small birds and small mammals appear to be the most frequently taken prey.

Short-eared Owl, a very rare species in Ireland, has been known to breed within the site. Nesting certainly occurred in the late 1970s and birds have been recorded intermittently since. The owls are considered to favour this site due to the presence of Bank Voles, a favoured prey item. Merlin also breed within the site but the size of the population is not known. Red Grouse is found on some of the unplanted areas of bog and heath – this is a species that has declined in Ireland and is now Red-listed.

The Stack's to Mullaghareirk Mountains, West Limerick Hills and Mount Eagle SPA is of ornithological importance because it provides excellent nesting and foraging habitat for breeding Hen Harrier and is one the top sites in the country for the species. The presence of three species, Hen Harrier, Merlin and Short-eared Owl, which are listed on Annex I of the E.U. Birds Directive is of note.

*For inspection purposes only.
Consent of copyright owner required for any other use.*

20.1.2015

SITE SYNOPSIS

SITE NAME: RIVER SHANNON AND RIVER FERGUS ESTUARIES SPA

SITE CODE: 004077

The estuaries of the River Shannon and River Fergus form the largest estuarine complex in Ireland. The site comprises the entire estuarine habitat from Limerick City westwards as far as Doonaha in Co. Clare and Dooneen Point in Co. Kerry.

The site has vast expanses of intertidal flats which contain a diverse macro-invertebrate community, e.g. *Macoma-Scrobicularia-Nereis*, which provides a rich food resource for the wintering birds. Salt marsh vegetation frequently fringes the mudflats and this provides important high tide roost areas for the wintering birds. Elsewhere in the site the shoreline comprises stony or shingle beaches.

The site is a Special Protection Area (SPA) under the E.U. Birds Directive, of special conservation interest for the following species: Cormorant, Whooper Swan, Light-bellied Brent Goose, Shelduck, Wigeon, Teal, Pintail, Shoveler, Scaup, Ringed Plover, Golden Plover, Grey Plover, Lapwing, Knot, Dunlin, Black-tailed Godwit, Bar-tailed Godwit, Curlew, Redshank, Greenshank and Black-headed Gull. It is also of special conservation interest for holding an assemblage of over 20,000 wintering waterbirds. The E.U. Birds Directive pays particular attention to wetlands and, as these form part of this SPA, the site and its associated waterbirds are of special conservation interest for Wetland & Waterbirds.

The site is the most important coastal wetland site in the country and regularly supports in excess of 50,000 wintering waterfowl (57,133 - five year mean for the period 1995/96 to 1999/2000), a concentration easily of international importance. The site has internationally important populations of Light-bellied Brent Goose (494), Dunlin (15,131), Black-tailed Godwit (2,035) and Redshank (2,645). A further 17 species have populations of national importance, i.e. Cormorant (245), Whooper Swan (118), Shelduck (1,025), Wigeon (3,761), Teal (2,260), Pintail (62), Shoveler (107), Scaup (102), Ringed Plover (223), Golden Plover (5,664), Grey Plover (558), Lapwing (15,126), Knot (2,015), Bar-tailed Godwit (460), Curlew (2,396), Greenshank (61) and Black-headed Gull (2,681) - figures are five year mean peak counts for the period 1995/96 to 1999/2000. The site is among the most important in the country for several of these species, notably Dunlin (13 % of national total), Lapwing (6% of national total) and Redshank (9% of national total).

The site also supports a nationally important breeding population of Cormorant (93 pairs in 2010).

Other species that occur include Mute Swan (103), Mallard (441), Red-breasted Merganser (20), Great Crested Grebe (50), Grey Heron (38), Oystercatcher (551),

Turnstone (124) and Common Gull (445) - figures are five year mean peak counts for the period 1995/96 to 1999/2000.

Apart from the wintering birds, large numbers of some species also pass through the site whilst on migration in spring and/or autumn.

The River Shannon and River Fergus Estuaries SPA is an internationally important site that supports an assemblage of over 20,000 wintering waterbirds. It holds internationally important populations of four species, i.e. Light-bellied Brent Goose, Dunlin, Black-tailed Godwit and Redshank. In addition, there are 17 species that have wintering populations of national importance. The site also supports a nationally important breeding population of Cormorant. Of particular note is that three of the species which occur regularly are listed on Annex I of the E.U. Birds Directive, i.e. Whooper Swan, Golden Plover and Bar-tailed Godwit. Parts of the River Shannon and River Fergus Estuaries SPA are Wildfowl Sanctuaries.

*For inspection purposes only.
Consent of copyright owner required for any other use.*

30.5.2015

*For inspection purposes only.
Consent of copyright owner required for any other use.*



Site Name: Moanveanlagh Bog SAC

Site Code: 002351

Moanveanlagh Bog is situated in Co. Kerry approximately 6 km east of Listowel, mainly within the townlands of Carhooeara and Bunagarha. The site comprises a raised bog that includes both areas of high bog and cutover bog.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes):

[7110] Raised Bog (Active)*
[7120] Degraded Raised Bog
[7150] Rhynchosporion Vegetation

Active raised bog comprises areas of high bog that are wet and actively peat-forming, where the percentage cover of bog mosses (*Sphagnum* spp.) is high, and where some or all of the following features occur: hummocks, pools, wet flats, *Sphagnum* lawns, flushes and soaks. Degraded raised bog corresponds to those areas of high bog whose hydrology has been adversely affected by peat cutting, drainage and other land use activities, but which are capable of regeneration. The Rhynchosporion habitat occurs in wet depressions, pool edges and erosion channels where the vegetation includes White Beak-sedge (*Rhynchospora alba*) and/or Brown Beak-sedge (*R. fusca*), and at least some of the following associated species, Bog Asphodel (*Narthecium ossifragum*), sundews (*Drosera* spp.), Deergrass (*Scirpus cespitosus*) and Carnation Sedge (*Carex panicea*).

This is a relatively flat site with some marginal areas that slope relatively steeply towards the cutover. There are a few large hummocks but over much of the site the micro-topography is very uniform. A flush area extends along the north and north-east of the site. In the south-west a bog burst has occurred and concentrically arranged tear pools can be seen, some of which are up to 12 m long. A swallow hole occurs near the middle of the site. Cutover bog occurs around the south-west, south and south-eastern margins of the high bog.

Much of the high bog has vegetation typical of a Western Raised Bog. The vegetation of the high bog is dominated by Bog Asphodel, White Beak-sedge, Cross-leaved Heath (*Erica tetralix*) and Carnation Sedge. Small patches of the moss *Racomitrium lanuginosum* and Common Lousewort (*Pedicularis sylvatica*) occur at the site. Purple Moor-grass (*Molinia caerulea*) is very common in the flush areas. The tear pools are mostly bare of vegetation but some support bladderwort (*Utricularia* sp.) and the bog mosses *S. cuspidatum* and *S. auriculatum*, with *S. papillosum* and the moss *Campylopus*

atrovirens occurring at the pool edges. Towards the margins of the bog Bog-myrtle (*Myrica gale*) is frequent.

Current land uses on the site consist of a small area of peat-cutting at the margins and a low level of grazing by cattle in the north-east section of the high bog. Peat-cutting has significantly declined since the 1970s. Other damaging operations include extensive fire damage, which is still occurring, and the dumping of household refuse and cars around the high bog. These are all activities that have resulted in the loss of habitat and damage to the hydrological status of the site, and pose a continuing threat to its viability. This site also suffers from invasive species, with the shrub *Rhododendron ponticum* recorded on the western edge of the site and the carnivorous Pitcher Plant (*Sarracenia purpurea*) forming a large colony.

Moanveanlagh Bog is significant in terms of its geographical location as it is at the extreme south-western range of raised bogs in Ireland. Moanveanlagh Bog is a site of considerable conservation significance as it comprises a raised bog, a rare habitat in the E.U. and one that is becoming increasingly scarce and under threat in Ireland. This site supports a good diversity of raised bog microhabitats, including flushes. Active raised bog is listed as a priority habitat on Annex I of the E.U. Habitats Directive. Priority status is given to habitats and species that are threatened throughout the E.U. Ireland has a high proportion of the total E.U. resource of this habitat type (over 60%) and so has a special responsibility for its conservation at an international level.

For inspection purposes only
Consent of copyright owner required for any other use



**CONSULTANTS IN ENGINEERING,
ENVIRONMENTAL SCIENCE
& PLANNING**

www.fehilytimoney.ie

For inspection purposes only.
Consent of copyright owner required for any other use.

CORK OFFICE

Core House
Pouladuff Road,
Cork, T12 D773,
Ireland
+353 21 496 4133

Dublin Office

J5 Plaza,
North Park Business Park,
North Road, Dublin 11, D11 PXT0,
Ireland
+353 1 658 3500

Carlow Office

Unit 6, Bagenalstown Industrial
Park, Royal Oak Road,
Muine Bheag,
Co. Carlow, R21 XW81,
Ireland
+353 59 972 3800

